



Basic Research in Computer Science

BRICS RS-97-16 Alon et al.: Linear Hashing

Linear Hashing

Noga Alon
Martin Dietzfelbinger
Peter Bro Miltersen
Erez Petrank
Gábor Tardos

BRICS Report Series

ISSN 0909-0878

RS-97-16

June 1997

**Copyright © 1997, BRICS, Department of Computer Science
University of Aarhus. All rights reserved.**

**Reproduction of all or part of this work
is permitted for educational or research use
on condition that this copyright notice is
included in any copy.**

**See back inner page for a list of recent BRICS Report Series publications.
Copies may be obtained by contacting:**

**BRICS
Department of Computer Science
University of Aarhus
Ny Munkegade, building 540
DK-8000 Aarhus C
Denmark
Telephone: +45 8942 3360
Telefax: +45 8942 3255
Internet: BRICS@brics.dk**

**BRICS publications are in general accessible through the World Wide
Web and anonymous FTP through these URLs:**

`http://www.brics.dk`
`ftp://ftp.brics.dk`
This document in subdirectory RS/97/16/

Linear Hashing

Noga Alon* Martin Dietzfelbinger† Peter Bro Miltersen‡
Erez Petrank§ Gábor Tardos¶

Abstract

Consider the set \mathcal{H} of all linear (or affine) transformations between two vector spaces over a finite field F . We study how good \mathcal{H} is as a class of hash functions, namely we consider hashing a set S of size n into a range having the same cardinality n by a randomly chosen function from \mathcal{H} and look at the expected size of the largest hash bucket. \mathcal{H} is a universal class of hash functions for any finite field, but with respect to our measure different fields behave differently.

*Dep. of Math., Sackler Faculty of Exact Sciences, Tel Aviv University, Tel Aviv, Israel and Institute for Advanced Study, Princeton, NJ 08540. Research supported in part by a USA-Israeli BSF grant, by the Sloan Foundation grant No. 96-6-2, by an NEC Research Institute grant and by the Hermann Minkowski Minerva Center for Geometry at Tel Aviv University. E-mail: noga@math.tau.ac.il.

†Fachbereich Informatik, Lehrstuhl II, Universität Dortmund, D-44221 Dortmund, Germany. E-mail: dietzf@ls2.informatik.uni-dortmund.de. Partially supported by DFG grant Di 412/5-1.

‡BRICS, Centre of the Danish National Research Foundation, University of Aarhus, Ny Munkegade, Aarhus, Denmark. Supported by the ESPRIT Long Term Research Programme of the EU under project number 20244 (ALCOM-IT). E-mail: bromille@brics.dk. Part of this work was done while the author was at the University of Toronto.

§DIMACS, P.O.Box 1179, Piscataway, NJ 08855-1179, USA. E-mail: erez@dimacs.rutgers.edu. Part of this work was done while the author was visiting the University of Toronto.

¶Mathematical Institute of the Hungarian Academy of Sciences, Pf. 127, Budapest, H-1364 Hungary. Supported by NSF grants CCR-95-03254 and DMS-9304580, a grant from Fuji Bank and the grant OTKA-F014919. E-mail: tardos@cs.elte.hu. Part of this work was done while the author was visiting the University of Toronto and the Institute for Advanced Study, Princeton.

If the finite field F has n elements then there is a bad set $S \subset F^2$ of size n with expected maximal bucket size $\Omega(n^{1/3})$. If n is a perfect square then there is even a bad set with largest bucket size *always* at least \sqrt{n} . (This is worst possible, since with respect to a universal class of hash functions every set of size n has expected largest bucket size below $\sqrt{n} + 1/2$.)

If, however, we consider the field of two elements then we get much better bounds. The best previously known upper bound on the expected size of the largest bucket for this class was $O(2\sqrt{\log n})$. We reduce this upper bound to $O(\log n \log \log n)$. Note that this is not far from the guarantee for a random function. There, the average largest bucket would be $\Theta(\log n / \log \log n)$.

In the course of our proof we develop a tool which may be of independent interest. Suppose we have a subset S of a vector space D over \mathbf{Z}_2 , and consider a random linear mapping of D to a smaller vector space R . If the cardinality of S is larger than $c_\epsilon |R| \log |R|$ then with probability $1 - \epsilon$, the image of S will cover all elements in the range.

1 Introduction

Consider distributing n balls in s buckets, randomly and independently. The resulting distribution of the balls in the buckets is the object of occupancy theory.

In the theory of algorithms and in complexity theory, it is often necessary and useful to consider putting balls in buckets without complete independence. More precisely, the following setting is studied: A class \mathcal{H} of hash functions, each mapping a universe U to $\{1, 2, \dots, s\}$, is fixed. A set $S \subseteq U$ to be hashed is given by an adversary, a member $h \in \mathcal{H}$ is chosen uniformly at random, S is hashed using h , and the distribution of the multi-set $\{h(x) | x \in S\}$ is studied. If the class \mathcal{H} is the class of all functions between U and $\{1, 2, \dots, s\}$, we get the classical occupancy problems. Carter and Wegman defined a class \mathcal{H} to be universal if¹

$$\forall x \neq y \in U : \text{Prob}(h(x) = h(y)) \leq 1/s.$$

¹We remark that a stricter definition is often used in the complexity theory literature, and a more liberal definition is often used in the data structure literature.

For universal families, the following properties are well known; variations of them have been used extensively in various settings:

1. If S of size n is hashed to n^2 buckets, with probability more than $1/2$, no collision occurs.
2. If S of size $2n^2$ is hashed to n buckets, with probability more than $1/2$, every bucket receives an element.
3. If S of size n is hashed to n buckets, the expected size of the largest bucket is less than $\sqrt{n} + \frac{1}{2}$.

The intuition behind universal hashing is that we often lose relatively little compared to using a completely random map. Note that for the property 1, this is true in a very strong sense; even with complete randomness, we do not expect $o(n^2)$ buckets to suffice (the birthday paradox), so nothing is lost by using a universal family instead. The bounds in the second and third properties, however, are rather coarse-grained compared to what one would get with complete randomness. For property 2, with complete randomness, $\Theta(n \log n)$ balls would suffice to cover the buckets with good probability (the coupon collector's theorem), i.e. a polynomial improvement over n^2 , and for property 3, with complete randomness, we expect the largest bucket to have size $\Theta(\log n / \log \log n)$, i.e. an exponential improvement over \sqrt{n} . In these last cases we do seem to lose quite a lot compared to using a completely random map and better bounds would seem desirable. However, it is rather easy to construct (unnatural) families of universal families and sets to be hashed showing that size $\Theta(n^2)$ is necessary to cover n buckets with non-zero probability, and that buckets of size \sqrt{n} are, in general unavoidable, when a set of size n is hashed to n buckets. This shows that the *abstract* property of universality does not allow for stronger statements. Now fix a *concrete* universal family of hash functions. We ask the following question: *To which extent are the fine-grained occupancy properties of completely random maps preserved?*

We provide answers to these questions for the case of *linear maps* between two vector spaces over a finite field, a natural and well known class of universal hash functions. The general flavor of our results is that “large fields are bad”, in the sense that the bounds becomes the worst possible for universal families, while “small fields are good” in the sense that the bounds

become as good or almost as good as the ones for independently distributed balls.

More precisely, for the covering problem, we show the following (easy) theorem

Theorem 1 *Let F be a field of size n and let \mathcal{H} be the class of linear maps between F^2 and F . There is a subset S of F^2 of size $\Theta(|F|^2)$, so that for no $h \in \mathcal{H}$, $h(S) = F$.*

On the other hand, we prove the following harder theorem

Theorem 2 *Let S be a subset of a vector space over \mathbf{Z}_2 and choose a random linear map to a smaller vector space R . If $|S| \geq c_\epsilon |R| \log |R|$ then with probability at least $1 - \epsilon$ the image of S covers the entire range R .*

For the “largest bucket problem”, let us first introduce some notation: Let U be the universe from which the keys are chosen. We fix a class \mathcal{H} mapping U to $\{1, \dots, s\}$. Then, a set $S \subseteq U$ of size n is chosen *by an adversary*, and we uniformly at random pick a hash function $h \in \mathcal{H}$, hash S using h and look at the size of the largest resulting hash bucket. We denote the expectation of this size by L_n^s . Formally,

$$L_n^s(\mathcal{H}) = \max_{S \subseteq U, |S|=n} E_{h \in \mathcal{H}} \left[\max_{y \in \{1, \dots, s\}} |\{x \in S \mid h(x) = y\}| \right]$$

Usually we think of s being of size close to n . Note that if $s = \Omega(n^2)$, any universal class yields $L_n^s = O(1)$.

The class \mathcal{H} we will consider is the set of linear maps between $F^m \rightarrow F^k$ for $m > k$. Here F is a finite field and $s = |F|^k$. This class is universal for all values of the parameters.

When $k = 1$ and thus $|F| = s$ the expected largest bucket can be large.

Theorem 3 *Let F be a finite field with $|F| = s$. For the class \mathcal{H} of all linear transformations $F^2 \rightarrow F$ we have*

$$L_s^s(\mathcal{H}) = \Omega(s^{1/3}).$$

Furthermore if $|F|$ is a perfect square we have

$$L_s^s(\mathcal{H}) > \sqrt{s}.$$

Note how close our lower bound for quadratic fields is to the upper bound of $\sqrt{s} + 1/2$ that holds for every universal class. We also mention that for the bad set we construct in Theorem 8 for a quadratic field there is no good hash function, since there *always* exists a bucket of size at least \sqrt{s} .

When the field is the field of two elements, the situation is completely different. Markowsky, Carter and Wegman [MCW78] showed that for this case $L_s^s(\mathcal{H}) = O(s^{1/4})$. Mehlhorn and Vishkin [MV84] improved on this result (although this is implicit in their paper) and showed that $L_s^s(\mathcal{H}) = O(2\sqrt{\log s})$. We further improve the bound and show that:

Theorem 4 *For the class \mathcal{H} of all linear transformations between two vector spaces over \mathbf{Z}_2 ,*

$$L_s^s(\mathcal{H}) = O(\log s \log \log s).$$

Furthermore, we also show that even if the range is smaller than $|S|$ by a logarithmic factor, the same still holds:

Theorem 5 *For the class \mathcal{H} of all linear transformations between two vector spaces over \mathbf{Z}_2 ,*

$$L_{s \log s}^s(\mathcal{H}) = O(\log s \log \log s).$$

Note that even if one uses the class \mathcal{R} of *all* functions one obtains only a slightly better result: the expected size of the largest bucket in this case is $L_s^s(\mathcal{R}) = \Theta(\log s / \log \log s)$ and $L_{s \log s}^s(\mathcal{R}) = \Theta(\log s)$, which is the best possible bound for any class. Interestingly, our upper bound is based on our upper bound for the covering property.

We do not know what the right bound is for the class of linear maps over \mathbf{Z}_2 , i.e., is it as good as $O(\log s / \log \log s)$? We leave this as an open question.

1.1 Motivation

There is no doubt that the method of implementing a dictionary by hashing with chaining, recommended in textbooks [CLR90, GBY90] especially for situations with many update operations, is a practically important scheme.

In situations in which a good bound on the cost of single operations is important, e. g., in real-time applications, the expected maximal bucket size as formed by all keys ever present in the dictionary during a time interval

plays a crucial role. Our results show that, at least as long as the size of the hash table can be determined right at the start, using a hash family of linear functions over \mathbf{Z}_2 will perform very well in this respect. For other simple hash classes such bounds on the worst case bucket size are not available or are even wrong (see Theorem 8); other, more sophisticated hash families [S89, DM90, DGMP92] that do guarantee small maximal bucket sizes consist of functions with higher evaluation time. Of course, if worst case constant time for certain operations is absolutely necessary, the known two-level hashing schemes can be used, e. g., the FKS scheme [FKS84] for static dictionaries; dynamic perfect hashing [DKMHRT94] for the dynamic case with constant time lookups and expected time $O(n)$ for n update operations; and the “real-time dictionaries” from [DM90] that perform each operation in constant time, with high probability. It should be noted, however, that a price is to be paid for the guaranteed constant lookup time in the dynamic schemes: the (average) cost of insertions is significantly higher than in simple schemes like chained hashing; the overall storage requirements are higher as well.

1.2 Related work

Another direction in trying to show that a specific class has a good bound on the expected size of the largest bucket is to build a class specifically designed to have such good property.

One immediate such result is obtained by looking at the class of d -degree polynomials over finite fields, where $d = c \log n / \log \log n$ (see, e.g., [ABI86].) It is easy to see that this class maps each d elements of the domain independently to the range, and thus, the bound that applies to the class of all functions also applies to this class. We can combine this with the following well known construction, (which is usually called “collapsing the universe”): At the expense of $\Theta(\log n + \log \log |U|)$ random bits one can construct a map $g : U \rightarrow \{1, \dots, n^{k+2}\}$ that on any set S will be one-to-one with probability $1 - O(1/n^k)$. This gives us a class with $L_n^n = \Theta(\log n / \log \log n)$ of size $2^{O(\log \log |U| + \log^2 n / \log \log n)}$ and with evaluation time $O(\log n / \log \log n)$ in a reasonable model of computation, say, a RAM with unit cost operations on members of the universe to be hashed.

More efficient (but much larger) families were given by Siegel [S89] and by Dietzfelbinger and Meyer auf der Heide [DM90]. Both provide families of size $|U|^{n^\epsilon}$ such that the functions can be evaluated in $O(1)$ time on a RAM

and with $L_n^n = \Theta(\log n / \log \log n)$. The families from [S89] and [DM90] are somewhat complex to implement while the class of linear maps requires only very basic bit operations (as discussed already in [CW79]). It is therefore desirable to study this class, and this is the main purpose of the present paper.

1.3 Notation

If S is a subset of the domain D of a function h we use $h(S)$ to denote $\{h(s) \mid s \in S\}$. If x is an element of the range we use $h^{-1}(x)$ to denote $\{s \in D \mid h(s) = x\}$. If A and B are subsets of a vector space V and $x \in V$ we use the notations $A+B = \{a+b \mid a \in A \wedge b \in B\}$ and $x+A = \{x+a \mid a \in A\}$. We use \mathbf{Z}_2 to denote the two element field. All logarithms in this paper are base two.

2 The covering property

2.1 Lower bounds for covering with a large field

We prove Theorem 1. Take a set $A \subset F$ of size $|A| = \lfloor |F|/2 \rfloor$ and consider $S = \{(x, y) \mid y \neq 0 \wedge x/y \in A \wedge (x-1)/y \notin A\}$. S has density around one quarter and no linear map $g : F^2 \rightarrow F$ satisfies $g(S) = F$. To see this take a nonzero linear map $g : (x, y) \mapsto ax + by$ and note that if $0 \in g(S)$ then $a \neq 0$ and $-b/a \in A$ but in this case $a \notin g(S)$.

2.2 Upper bounds for covering with a small field - the existential case

We start by showing that if we have a large enough subset A of a vector space over \mathbf{Z}_2 then *there exists* a linear transformation T to a large vector space such that $T(A)$ is the entire range. The constant e below is the base of the natural logarithm.

Theorem 6 *Let A be a finite set of vectors in a vector space V of an arbitrary dimension over \mathbf{Z}_2 and let $t > 0$ be an integer. If $|A| > t2^t / \log e$ then there exists a linear map $T : V \rightarrow \mathbf{Z}_2^t$, so that T maps A onto \mathbf{Z}_2^t .*

For the proof of this theorem we need the following simple lemma. Note that although we state the lemma for vector spaces, it holds for any finite group.

Lemma 2.1 *Let V be a finite vector space, $A \subseteq V$, $\alpha = 1 - |A|/|V|$. Then for a random $v \in V$ it holds that*

$$E_v(1 - |A \cup (v + A)|/|V|) = \alpha^2.$$

Proof. If v and u are both chosen uniformly independently at random from V then both events $u \notin A$ and $u \notin v + A$ have probability α and they are independent. \square

Proof of Theorem 6. Let m be the dimension of V , $N = |A|$ and $\alpha = 1 - |A|/|V| = 1 - N/2^m$. Starting with $A_0 = A$, we choose a vector $v_1 \in V$ so that for $A_1 = A_0 \cup (v_1 + A_0)$

$$1 - \frac{|A_1|}{|V|} \leq \alpha^2.$$

Such a choice for v_1 exists by Lemma 2.1. Then, by the same procedure, we choose a v_2 so that for

$$A_2 = A_1 \cup (v_2 + A_1) = A + \text{Span}\{v_1, v_2\},$$

$$1 - \frac{|A_2|}{|V|} \leq \alpha^4,$$

and so on up to $A_s = A + \text{Span}\{v_1, \dots, v_s\}$ with $s = m - t$ for which

$$1 - \frac{|A_s|}{|V|} \leq \alpha^{2^s}.$$

Note that one can assume that the vectors v_1, \dots, v_s are linearly independent since choosing a vector v_i which depends on the previous choices makes $A_i = A_{i-1}$.

We now claim that $A_s = V$. Suppose in way of contradiction that $A_s \neq V$. Take a vector $v \in V$ outside A_s , then $v + \text{Span}\{v_1, \dots, v_s\}$ is disjoint from A_s ; thus

$$1 - \frac{|A_s|}{|V|} \geq 2^s/2^m.$$

Combining the last two inequalities and recalling that $t = m - s$ we get

$$2^{-t} \leq \alpha^{2^s} = (1 - N/2^m)^{2^s} < e^{-N2^{-t}}.$$

Taking the logarithm of both sides yields $-t < -N \log_2 e / 2^t$ contradicting our assumption on N . Thus, $A_s = V$ must hold.

Now we choose a linear map $T : V \rightarrow \mathbf{Z}_2^t$ such that its kernel $T^{-1}(0) = \text{Span}\{v_1, \dots, v_s\}$. The equality $V = A_s = A + \text{Span}\{v_1, \dots, v_s\}$ implies that T maps A onto \mathbf{Z}_2^t . \square

The bound in Theorem 6 is asymptotically tight as shown by the following proposition.

Proposition 2.2 *For every large enough integer t there is a set A of at least $(t - 3 \log t)2^t / \log e$ vectors in a vector space V over \mathbf{Z}_2 so that for any linear map $T : V \rightarrow \mathbf{Z}_2^t$, T does not map A onto \mathbf{Z}_2^t .*

Proof: Let $V = \mathbf{Z}_2^{t+s}$ with $s = \lfloor t/10 \rfloor$ and let A be chosen at random by picking each element of V independently and with probability $p = 1 - 2^{-x}$ into the set with $x = (t - 2 \log t)2^{-s}$. From Chebyshev's inequality we know that with probability at least $3/4$, A has cardinality at least $pN - 2\sqrt{pN}$ for $N = 2^{s+t}$. Using $p > x / \log e - x^2 / (2 \log^2 e)$ one can show that this is as many as claimed in the proposition. Let us compute the probability that there exists a linear map $T : V \rightarrow \mathbf{Z}_2^t$ such that T maps A onto \mathbf{Z}_2^t . There are $2^{t(t+s)}$ possible maps T and each of them satisfies $T(A) = \mathbf{Z}_2^t$ with probability at most $(1 - (1 - p)^{2^s})^{2^t} = (1 - 2^{-2^s x})^{2^t} = (1 - t^2 / 2^t)^{2^t} < e^{-t^2}$. So with probability almost $3/4$, A is not small and still no T maps A onto \mathbf{Z}_2^t . \square

2.3 Choosing the linear map at random

In this subsection we strengthen Theorem 6 and prove that if A is bigger than what is required there by only a constant factor, then almost all choices of the linear transformation T work. This may seem immediate at first glance since Lemma 2.1 tells us that a random choice for the next vector is good on average. In particular, it might seem that for a random choice of v_1 and v_2 in the proof of Theorem 6, $E_{v_1, v_2}(1 - |A + \text{Span}\{v_1, v_2\}| / |V|) \leq \alpha^4$. Unfortunately this is not the case: For example, think of A being a linear

subspace containing half of V . In this case, the ratio α of points that are not covered is $1/2$. As random vectors v_i are chosen to be added to A , vectors in A are chosen with probability $1/2$. Thus, after i steps, α remains $1/2$ with probability $1/2^i$ and becomes 0 otherwise. Thus, the expected value of α_i is 2^{-i-1} which is much bigger than 2^{-2^i} .

Our first lemma is technical in nature.

Lemma 2.3 *Let α_i for $1 \leq i \leq k$ be random variables and let $0 < \alpha_0 < 1$ be a constant. Suppose that for $0 \leq i < k$ we have $0 \leq \alpha_{i+1} \leq \alpha_i$ and conditioned on any set of values for $\alpha_1, \dots, \alpha_i$ we have $E[\alpha_{i+1} | \alpha_1, \dots, \alpha_i] = \alpha_i^2$. Then for any threshold $0 < t < 1$ we have*

$$\text{Prob}[\alpha_k \geq t] \leq \alpha_0^{k - \log \log(1/t) + \log \log(1/\alpha_0)}.$$

Proof: The proof is by induction on k . The $k = 0$ base case is trivial.

We assume the statement of the lemma is true for k and prove it for $k+1$. Let $c = k - \log \log(1/t)$. We may suppose $c + 1 + \log \log(1/\alpha_0) \geq 0$ since otherwise the bound in the lemma is above 1.

After the choice of α_1 , the rest of the random variables form a random process of length k satisfying the conditions of the lemma (unless $\alpha_1 = 0$); thus we can apply the inductive hypothesis to get

$$\text{Prob}[\alpha_{k+1} \geq t] = E_{\alpha_1}[\text{Prob}[\alpha_{k+1} \geq t | \alpha_1]] \leq E[f(\alpha_1)],$$

where we define $f_0(x) = x^{c + \log \log(1/x)}$ for $0 < x < 1$ and take $f(x) = \min(1, f_0(x))$ in the same interval and $f(0) = 0$. The value $f(\alpha_1)$ is clearly an upper bound on $\text{Prob}[\alpha_{k+1} \geq t | \alpha_1]$.

We claim that in the interval $0 \leq x \leq \alpha_0$ we have $f(x) \leq f_0(\alpha_0)x/\alpha_0$. To prove this simply observe that $f_0(x)/x$ is first increasing then decreasing on $(0, 1)$. To see this compute the derivative $(f_0(x)/x)' = (c + \log e - 1 + \log \log(1/x))f_0(x)/x^2$. If α_0 is still in the increasing phase then we have $f(x)/x \leq f_0(x)/x \leq f_0(\alpha_0)/\alpha_0$ for $0 < x \leq \alpha_0$. Suppose now that α_0 is already in the decreasing phase and define $x' = 2^{-2^{-c-1}}$. Notice that we assumed $\alpha_0 \leq x'$ in the beginning of the proof, so we have $f_0(\alpha_0)/\alpha_0 \geq f_0(x')/x'$. Let us define $x'' = x'^2 = 2^{-2^{-c}}$ and notice that we have $f(x) = 1$ if and only if $x \geq x''$. It is easy to check that x'' must still be in the increasing phase of $f_0(x)/x$ thus we have $f(x)/x = f_0(x)/x \leq f_0(x'')/x'' = 1/x''$ for $0 < x \leq x''$. For $x'' \leq x < 1$ we simply have $f(x)/x = 1/x \leq 1/x''$. Thus we

must have $f(x)/x \leq 1/x'' = f_0(x')/x' \leq f_0(\alpha_0)/\alpha_0$ for $0 < x < 1$. We have thus proved the claim in all cases for $0 < x \leq \alpha_0$. The claim is trivial for $x = 0$.

Using the claim we can finish the proof writing:

$$\text{Prob}[\alpha_{k+1} \geq t] \leq E[f(\alpha_1)] \leq E[f_0(\alpha_0)\alpha_1/\alpha_0] = f_0(\alpha_0)E[\alpha_1]/\alpha_0 =$$

$$f_0(\alpha_0)\alpha_0 = \alpha_0^{c+1+\log \log(1/\alpha_0)}.$$

□

We remark that the bound in the lemma is achievable for $t = \alpha_0^{2^j}$ with an integer $0 \leq j \leq k$. The optimal process has $\alpha_i = \alpha_{i-1}$ or $\alpha_i = 0$ for $1 \leq i \leq k - j$, while $\alpha_i = \alpha_{i-1}^2$ for $k - j < i \leq k$.

Theorem 7 *a) For every $\epsilon > 0$ there is a constant $c_\epsilon > 0$ such that the following holds. Let A be a finite set of vectors in a vector space V of an arbitrary dimension over \mathbf{Z}_2 , let $t > 0$ be an integer. If $|A| \geq c_\epsilon t^t$ then for a uniform random linear transformation $T : V \rightarrow \mathbf{Z}_2^t$*

$$\text{Prob}(T(A) = \mathbf{Z}_2^t) \geq 1 - \epsilon.$$

b) If A is a subset of the vector space \mathbf{Z}_2^u of density $|A|/2^u = 1 - \alpha < 1$ and $0 \leq t \leq u$ is an integer then for a uniform random onto linear transformation $T : \mathbf{Z}_2^u \rightarrow \mathbf{Z}_2^t$

$$\text{Prob}(T(A) \neq \mathbf{Z}_2^t) \leq \alpha^{u-t-\log t+\log \log(1/\alpha)}.$$

Proof: We start with proving part b) of the theorem. In order to pick the onto map T we use the following process (similar to the one in the proof of Theorem 6). Pick $s = u - t$ vectors v_1, \dots, v_s uniformly at random from the vectors in \mathbf{Z}_2^u and choose T to be a random onto linear transformation $T : \mathbf{Z}_2^u \rightarrow \mathbf{Z}_2^t$ with the constraints $T(v_i) = 0$ ($i = 1, \dots, s$), i.e. the vectors v_1, \dots, v_s are in the kernel of T . Note that the v_i 's are not necessarily linearly independent and that they do not necessarily span the kernel. After picking them, an *onto* transformation satisfying that the v_i 's are in the kernel is chosen at random.

Using notations similar to the ones used in the proof of Theorem 6, let $A_0 = A$, $A_i = A_0 + \text{Span}\{v_1, \dots, v_i\}$ and $\alpha_i = 1 - |A_i|/2^u$ for $i = 0, \dots, s$. Clearly α_i is nonnegative and monotone decreasing in i with $\alpha_0 = \alpha$. The

equation $E[\alpha_{i+1} \mid \alpha_1, \dots, \alpha_i] = \alpha_i^2$ is guaranteed by Lemma 2.1 since $A_{i+1} = A_i + v_{i+1}$ and v_{i+1} is independent of α_j for $j \leq i$. Thus all the conditions of Lemma 2.3 are satisfied and we have

$$\text{Prob}[\alpha_s \geq 2^{-t}] \leq \alpha^{s - \log t + \log \log(1/\alpha)}.$$

By the definition of s the right hand side here is equal to the estimate in the theorem. Finally notice that if $T(A) \neq \mathbf{Z}_2^t$ then for the vector $w \in \mathbf{Z}_2^t$ not in $T(A)$ the set $T^{-1}(w)$ of size 2^{u-t} is disjoint from A_s and thus $\alpha_s = 1 - |A_s|/2^u \geq |T^{-1}(w)|/2^u = 2^{-t}$. Thus we have the claimed upper bound for the probability that $T(A) \neq \mathbf{Z}_2^t$.

Now we turn to part a) of the theorem and prove it using part b). Part a) is about a random linear transformation, not necessarily onto, but this difference from the claim just proved poses less of a problem, the difficulty is that we do not have an a priori bound on $1 - |A|/|V|$. In fact, this ratio can be arbitrarily small. To solve this, we choose the transformation T in two steps, the first step ensuring that the density of the covered set is substantial, then applying part b) for the second step.

Let $W = \mathbf{Z}_2^u$, with $u = \lceil \log(2|A|/\epsilon) \rceil$. We factor T through W . First, we pick uniformly at random a linear transformation $T_0 : V \rightarrow W$. Then, we pick a random onto linear map $T_1 : W \rightarrow \mathbf{Z}_2^t$, and set $T = T_0 \circ T_1$. This results in a uniformly chosen linear map $T : V \rightarrow \mathbf{Z}_2^t$.

Any pair of vectors $v \neq w \in A$ collide with probability $\text{Prob}[T_0(v) = T_0(w)] = 1/|W|$. Thus the expected number of collisions is $\binom{|A|}{2}/|W|$. Since $|T_0(A)| \leq |A|/2$ implies at least $|A|/2$ such collisions Markov's inequality gives $\text{Prob}[|T_0(A)| \leq |A|/2] \leq 2 \binom{|A|}{2} / (|A||W|) < |A|/|W| \leq \epsilon/2$. For any fixed T_0 part b) of the theorem gives

$$\text{Prob}[T(A) \neq \mathbf{Z}_2^t] \leq \alpha^{u-t-\log t + \log \log(1/\alpha)},$$

where $\alpha = 1 - |T_0(A)|/|W|$. In case $|T_0(A)| > |A|/2$ we have $\alpha < 1 - |A|/(2|W|) < e^{-\epsilon/8}$, thus using the monotonicity of the bound above we get

$$\text{Prob}[T(A) \neq \mathbf{Z}_2^t] \leq e^{-\epsilon(u-t-\log t + \log(\log e^{\epsilon/8})/8)}. \quad (1)$$

Choosing $c_\epsilon = 4(2/\epsilon)^{8/\epsilon}$ we have $|A| \geq c_\epsilon t 2^t$ and thus $u = \lceil \log(2|A|/\epsilon) \rceil > t + \log t + \log(8/\epsilon) + (8/\epsilon) \log(2/\epsilon)$. This implies that the bound in Equation 1

is below $\epsilon/2$, thus we get $\text{Prob}[T(A) \neq \mathbf{Z}_2^t] \leq \text{Prob}[|T_0(A)| \leq |A|/2] + \epsilon/2 < \epsilon$ as claimed. \square

We remark that a more careful analysis gives c_ϵ that is a small polynomial of $1/\epsilon$.

3 The largest bucket

3.1 Lower bound for the largest bucket with a large field

We start by showing why linear hashing over a large finite field is bad with respect to the expected largest bucket size measure. This natural example shows that universality of the class is not enough to assure small buckets. For a finite field F we prove the existence of a bad set $S \subset F^2$ of size $|S| = |F|$ such that the expected largest bucket in S with respect to a random linear map $F^2 \rightarrow F$ is big. We prove the results in Theorem 3 separately for quadratic and non-quadratic fields.

We start with an intuitive description of the constructions. Linear hashing of the plane collapses all straight lines of a random direction. Thus, a bad set in the plane must contain many points on at least one line in many different directions. It is not hard to come up with bad sets that contain many points of many different lines, however the obvious constructions (subplane or grid) yield sets where many of the ‘‘popular lines’’ tend to be parallel and thus they only cover a few directions. This problem can be solved by a projective transformation: the transformed set has many popular lines, but they are no longer parallel.

Theorem 8 *Let F be a finite field with $|F|$ being a perfect square. There exists a set $S \subset F^2$ of size $|S| = |F|$ such that for every linear map $h : F^2 \rightarrow F$, S has a large bucket, i.e. there exists a value $y \in F$ with $|h^{-1}(y)| \geq \sqrt{|F|}$.*

Proof. We have a finite field F_0 of which F is a quadratic extension. Let $|F_0| = m$ and $|F| = m^2 = n$. Let a be an arbitrary element in $F \setminus F_0$ and define $S = \{(\frac{1}{x+a}, \frac{y}{x+a}) \mid x, y \in F_0\}$. Note that $|S| = m^2 = |F|$. Notice also, that S is the image of the subplane F_0^2 under the projective transformation $(x, y) \mapsto (\frac{1}{x+a}, \frac{y}{x+a})$.

Fix $A, B \in F$ and consider the function $h : F^2 \rightarrow F$ defined by $h(x, y) = Ax + By$. We must show that there is some $C \in F$ such that $|h^{-1}(C) \cap S| \geq m$. If $B = 0$ then h maps all the m elements of $S' = \{(1/a, y/a) \mid y \in F_0\}$ to $C = A/a$, as needed. Otherwise, we claim that there is a $C \in F$ such that both $\frac{C}{B}$ and $\frac{aC-A}{B}$ are in F_0 . To see this observe that if g_1 and g_2 are two distinct members of F_0 , then ag_1 and ag_2 lie in distinct additive cosets of F_0 in F , since otherwise their difference, $a(g_1 - g_2)$ would have to be in F_0 , contradicting the fact that $a \notin F_0$. Thus, as g ranges over all members of F_0 , ag intersects distinct additive cosets of F_0 in F , and hence aF_0 intersects all those cosets. In particular, there is some $g \in F_0$ so that $ag \in F_0 + \frac{A}{B}$, implying that $C = gB$ satisfies the assertion of the claim. For the above C , for every choice of $x \in F_0$, $y(x) = \frac{C}{B}x + \frac{aC-A}{B} \in F_0$. We have now $A\frac{1}{a+x} + B\frac{y(x)}{a+x} = C$, showing that h maps all the m elements of $S' = \{(\frac{1}{a+x}, \frac{y(x)}{a+x}) \mid x \in F_0\} \subset S$ to C . \square

Theorem 9 *Let F be a finite field. There exists a set $S \subset F^2$ of size $|S| = |F|$ such that for more than half of the linear maps $h : F^2 \rightarrow F$, S has a large bucket, i.e. there exists a value $y \in F$ with $|h^{-1}(y)| \geq |F|^{1/3}/3 - 1$.*

Proof. First we construct a set $S' \subset F^2$ such that $|S'| \leq |F| = n$ and there are n distinct lines in the plane F^2 each containing at least $m \geq n^{1/3}/3$ points of S' .

Let us first consider the case when n is a prime, so F consists of the integers modulo n . We let $A = \{i \mid 1 \leq i < \sqrt{n}\}$ and consider the square grid $S' = A \times A$. Clearly $|S'| < n$. It is well known that each of the n most popular lines contains at least $m \geq n^{1/3}/3$ points of S' . This is usually proved for the same grid in the Euclidean plane (see e.g. [PA95], pp. 178–179) but that result implies the same for our grid in F^2 .

Now let $n = p^k$ and let F_0 be the subfield in F of p elements. Let $x \in F$ be a primitive element, then every element of F can be uniquely expressed as a polynomial of x of degree below k with coefficients from F_0 . Let $k_1 = \lfloor \frac{k+1}{3} \rfloor$, $k_2 = k - k_1 = \lfloor \frac{2k+1}{3} \rfloor$ and let $A_1 = \{f(x) \mid \deg(f) < k_1\}$, $A_2 = \{f(x) \mid \deg(f) < k_2\}$ where the polynomials f have coefficients from F_0 . Finally we take $S' = A_1 \times A_2$. Clearly $|S'| = n$. For $a \in A_1$ and $b \in A_2$ we consider the line $L_{a,b} = \{(y, ay + b) \mid y \in F\}$ in F^2 . Notice that there are n such lines and we have $ay + b \in A_2$ whenever $y \in A_1$. Thus, we have n distinct lines each containing $m = |A_1| = p^{k_1}$ points of S' . We have $m \geq n^{1/3}$

as claimed unless $k \equiv 1 \pmod{3}$. Notice that for $k \equiv 2 \pmod{3}$ our m is much higher than $n^{1/3}$. For the bad case $k \equiv 1 \pmod{3}$ we apply the construction below instead.

Finally suppose $n = p^k$, p is a prime and $k \equiv 1 \pmod{3}$. To get our set S' in this case we have to merge the two constructions above. Let F_0 be the p element subfield of F , then F_0 consists of the integers modulo p . We set $A = \{i \mid 1 \leq i < \sqrt{p}\}$. Let $k_1 = (k+2)/3$ and $k_2 = (2k+1)/3$ and let $x \in F$ be a primitive element, so we can express any element of F uniquely as a polynomial of x of degree less than k with coefficients from F_0 . We set $A_1 = \{f(x) \mid \deg(f) < k_1 \wedge f(0) \in A\}$, $A_2 = \{f(x) \mid \deg(f) < k_2 \wedge f(0) \in A\}$ where the polynomials f have coefficients from F_0 . Finally we set $S' = A_1 \times A_2$. Clearly $|S'| < n$. For $j, j' \in F_0$ let $L_{j,j'} = \{(i, ji + j') \mid i \in F_0\}$. Let a and b be polynomials with coefficients from F_0 with $\deg(a) < k_1$ and $\deg(b) < k_2$. Consider the line $L_{a,b} = \{(y, a(x)y + b(x)) \mid y \in F\}$. Notice that $|L_{a,b} \cap S'| = p^{k_1-1} |L_{a(0),b(0)} \cap (A \times A)|$. Thus, from knowing that the p most popular lines in F_0^2 contain at least $m_0 \geq p^{1/3}/3$ points from $A \times A$ we conclude that there exist n distinct lines each containing at least $m = m_0 p^{k_1-1} \geq n^{1/3}/3$ points of S' .

In all cases now we have constructed our set $S' \subset F^2$ of size $|S'| \leq n$ with n distinct popular lines each containing at least $m > n^{1/3}/3$ points of S' . Let P be the projective plane containing F^2 . Out of the $n^2 + n + 1$ points in P every popular line covers $n + 1$. The i th popular line ($1 \leq i \leq n$) can only have $i - 1$ intersections with earlier lines, thus it covers at least $n + 2 - i$ points previously uncovered. Therefore a total of at least $\binom{n+2}{2} - 1$ points are covered by popular lines. Simple counting gives the existence of a line L in P not among the popular lines, such that more than half of the points on L are covered by popular lines. Let f be a projective transformation taking the ideal line $L' = P \setminus F^2$ to L . We define $S = \{x \in F^2 \mid f(x) \in S'\}$. Clearly $|S| \leq |S'| \leq n$.

One linear hash function $h : F^2 \rightarrow F$ is constant zero (and thus all of S is a single bucket), for the rest there is a point $x_h \in L'$ such that h collapses the points of F^2 of each single line going through x_h . As we get all points of L' equal number of times ($n - 1$ times to be precise) it is enough to prove that whenever $f(x_h)$ is covered by a popular line S has a big bucket with respect to h as claimed. So suppose $f(x_h)$ is in the popular line L'' . As L'' contains at least m points of S' thus $f^{-1}(L'')$ must be a line through x_h containing at least $m - 1$ points of S (the -1 comes from the possibility of $f(x_h) \in S'$).

This ensures a bucket of S with respect to h of size at least $m - 1$ as claimed. \square

3.2 Upper bound for the largest bucket with a small field

Let us now recall and prove our main result.

For convenience here we speak about hashing $n \log n$ keys to n values. Also, we assume that n is a power of 2.

Theorem 5: Let \mathcal{H} be the class of linear transformations between two vector spaces over \mathbf{Z}_2 , then

$$L_{n \log n}^n(\mathcal{H}) = O(\log n \log \log n).$$

This theorem implies Theorem 4.

We have to bound the probability of the event that many elements in the set S are mapped to a single element in the range. Denote this bad event by E_1 . The overall idea is to present another (less natural) event E_2 and show that the probability of E_2 is small, yet the probability of E_2 given E_1 is big. Thus, the probability of E_1 must be small. We remark here that a somewhat similar line of reasoning was used in the seminal paper of Vapnik and Chervonenkis [VC71].

For the proof we fix the domain to be $D = \mathbf{Z}_2^m$, the range (the buckets) to be $B = \mathbf{Z}_2^{\log n}$, and $S \subset D$ of size $|S| = n \log n$.

Let us choose arbitrary $\ell \geq \log n$ and consider the space $A = \mathbf{Z}_2^\ell$. We construct the linear transformation $h : D \rightarrow B$ through the intermediate range A in the following way. We choose uniformly at random a linear transformation $h_1 : D \rightarrow A$ and uniformly at random an onto linear transformation $h_2 : A \rightarrow B$. Now we define $h \stackrel{\text{def}}{=} h_1 \circ h_2$. Note that as mentioned in the proof of part a) of Theorem 7 this yields an h which is uniformly chosen from among all linear transformations from D to B .

Let us fix a threshold t . We define two events. E_1 is the existence of a bucket of size at least t :

Event E_1 : There exists an element $\alpha \in B$ such that

$$|h^{-1}(\alpha) \cap S| > t.$$

We are going to limit the probability of E_1 through the seemingly unrelated event E_2 :

Event E_2 : There exists an element $\alpha \in B$ such that

$$h_2^{-1}(\alpha) \subseteq h_1(S).$$

Consider the distribution space in which h_1 and h_2 are uniformly chosen as above. We shall show that

Proposition 3.1 *If $d = 2^\ell / (n \log n) > 1$ we have*

$$\text{Prob}[E_2] \leq d^{-\log d - \log \log d}.$$

Proposition 3.2 *If $t > c_{1/2}(2^\ell/n) \log(2^\ell/n)$ (with $c_{1/2}$ from Theorem 7a)) then*

$$\text{Prob}[E_2|E_1] \geq \frac{1}{2}.$$

From Propositions 3.1 and 3.2 we deduce that the probability of E_1 must be small:

Corollary 3.3 *For every $c > 1$ and every power n of 2 a random linear transformation hashing a subset S of a \mathbf{Z}_2 vector space of size $|S| = n \log n$ to $\mathbf{Z}_2^{\log n}$ we have*

$$\text{Prob}[\text{maximum bucket size} \geq c \log n \log \log n] < c^{-(1-o(1)) \log c}.$$

Here the $o(1)$ term depends only on c .

Proof: We may suppose that c and n are sufficiently large. We set $c' = c/c_{1/2}$ with the absolute constant $c_{1/2}$ guaranteed by Theorem 7a) and let $\ell = \lceil \log n + \log \log n + \log c' - 2 \log c' / \log \log n \rceil$ to satisfy the condition of Proposition 3.2 with $t = c \log n \log \log n$. We get $\text{Prob}[E_2|E_1] \geq \frac{1}{2}$. Then applying 3.1 we get $d > c'^{1-2/\log \log n} / 2 = c^{1-o(1)}$, so in particular $d > 1$ if c is large enough. Thus, we have $\text{Prob}[E_1] \leq 2\text{Prob}[E_2] < 2d^{-\log d - \log \log d} = c^{-(1-o(1)) \log c}$. \square

Let us now prove the propositions above.

Proof of Proposition 3.1: Note first that an alternative way to describe E_2 is

$$h_2(A \setminus h_1(S)) \neq B.$$

We will prove that Proposition 3.1 holds for any specific h_1 , and thus it also holds for a randomly chosen h_1 . So fix h_1 and consider the distribution in which h_2 is chosen uniformly amongst all full rank linear transformation from A to B .

We use part b) of Theorem 7 for the set $A \setminus h_1(S) \subset A$. Its density is clearly $1 - \alpha$ for $\alpha = |h_1(S)|/|A| \leq |S|/|A| = 1/d$. Thus the theorem gives $\text{Prob}[E_2] \leq \alpha^{\ell - \log n - \log \log n + \log \log(1/\alpha)} \leq d^{-\log d - \log \log d}$ as claimed. \square

Proof of Proposition 3.2: Fix h for which E_1 holds, and fix any full rank h_2 . We will show that the probability of event E_2 is at least $1/2$ even when these two are fixed and thus the conditional probability is also at least $1/2$.

Now since E_1 holds there is a subset $S' \subseteq S$ of cardinality at least t mapped by h to a single element $\alpha \in \mathbf{Z}_2^{\log n}$. Fix this α and define $D' \stackrel{\text{def}}{=} h^{-1}(\alpha)$ and $A' \stackrel{\text{def}}{=} h_2^{-1}(\alpha)$. Consider the distribution of h_1 satisfying $h = h_1 \circ h_2$. When we restrict h_1 to D' , we get that the distribution implied by such h_1 is a uniform choice of an affine or linear map from D' into A' (we show this in Proposition 3.4 below). For event E_2 to hold it is enough to have $A' \subseteq h_1(S)$. We will show that $h_1(S')$ covers all the points in A' with probability at least $1/2$ and thus we get that event E_2 happens with probability $1/2$. Since h_2 is onto we have $|A'| = 2^\ell/n$. On the other hand, $D' \cap S$ has cardinality at least $t = \lceil c_{1/2}(2^\ell/n) \log(2^\ell/n) \rceil$. By part a) of Theorem 7, the probability that a set of cardinality t mapped by a random linear transformation will cover a range of cardinality $2^\ell/n$ is at least $1/2$. (Note that Theorem 7a) clearly applies to a random affine transformation too.) \square

At this point, we have proven Corollary 3.3. This limits the probability of large buckets with linear hashing. It is straightforward to deduce Theorem 5 from that corollary:

Proof of Theorem 5: $L_{n \log n}^n$ is the expectation of the distribution of the largest bucket size. Corollary 3.3 limits the probability of the tail of this distribution, thus yielding the desired bound on the expectation:

$$E[\max S\text{-bucket size}] = \int_0^\infty \text{Prob}[\max S\text{-bucket} > t] dt <$$

$$\log n \log \log n (c_0 + \int_{c_0}^{\infty} c^{-\log c/2} dc) = O(\log n \log \log n),$$

if the $o(1)$ term of Corollary 3.3 is below $1/2$ for $c > c_0$. \square

In order for the paper to be self-contained we include a proof of the simple statement about random linear transformations used above.

Proposition 3.4 *Let D , A and B be vector spaces over \mathbf{Z}_2 . Let $h : D \rightarrow B$ be an arbitrary linear map, and let $h_2 : A \rightarrow B$ be an arbitrary onto linear map. Let α be any point in B and denote $D' \stackrel{\text{def}}{=} h^{-1}(\alpha)$ and $A' \stackrel{\text{def}}{=} h_2^{-1}(\alpha)$. Then, choosing a uniform linear map $h_1 : D \rightarrow A$ such that $h = h_1 \circ h_2$ and restricting the domain to D' we get a uniformly chosen linear map from D' to A' if $\alpha = 0$ or uniformly chosen affine map from D' to A' otherwise.*

Proof: Consider $D_0 \stackrel{\text{def}}{=} h^{-1}(0)$ and $A_0 \stackrel{\text{def}}{=} h_2^{-1}(0)$. Let us choose a complement space D_1 to D_0 in D , i.e. $D_0 \cap D_1 = \{0\}$ and $D_0 + D_1 = D$. Let us call x the unique vector in $D' \cap D_1$. We have $D' = D_0 + x$. A linear transformation $h_1 : D \rightarrow A$ is determined by its two restrictions $h' : D_0 \rightarrow A$ and $h'' : D_1 \rightarrow A$. Clearly the uniform random choice of h_1 corresponds to uniform and independent choices for h' and h'' . The restriction $h = h_1 \circ h_2$ means that $h'(D_0) \subseteq A_0$ and $h'' \circ h_2$ is the restriction of h to D_1 . Thus, after the restriction the random choices of h' and h'' are still independent. Note now that if $\alpha = 0$ then the restriction of h_1 in question is exactly $h' : D' \rightarrow A'$. If $\alpha \neq 0$ then use $h_1(u + x) = h'(u) + h''(x)$ for $u \in D_0$ to note that the restriction in question is again h' , this time translated by the random value $h''(x) \in A'$. \square

4 Remarks and open questions

We have discussed the case of a very small field (size 2) and a very large field (size n). What happens with intermediate sized fields? Some immediate rough generalizations of our bounds are the following: If we hash an adversarially chosen subset of F^m of size $n = |F|^k$ to F^k by a randomly chosen linear map, the expected size of the largest bucket is at most $O((\log n \log \log n)^{\log |F|})$ and at least $\Omega(|F|^{1/3})$. Tighter bounds should be possible.

Another question is which fine-grained property other well known hash families have. Examples of the families we have in mind include: Arithmetic

over \mathbf{Z}_p [CW79, FKS84] (with $h_{a,b}(x) = (ax+b \bmod p) \bmod n$), integer multiplication [DHKP93, AHNR95] (with $h_a(x) = (ax \bmod 2^k) \operatorname{div} 2^{k-l}$), Boolean convolution [MNT93] (with $h_a(x) = a \circ x$ projected to some subspace).

An example of a natural non-linear scheme for which the assertion of Theorem 6 fails is the scheme that maps integers between 1 and p , for some large prime p , to integers between 0 and $n - 1$ for $n = \lceil p/m \rceil$, by mapping $x \in \mathbf{Z}_p$ to the integer part of the fraction $(ax + b) \bmod p / m$, where a, b are two randomly chosen elements of \mathbf{Z}_p . For this scheme, there are primes p and choices of an n and a subset S of cardinality $\Omega(n \log n \log \log \log n)$ of \mathbf{Z}_p , which is not mapped by the above mapping onto $[0, n - 1]$ under any choice of a and b .

To see this, let p be a prime satisfying $p \equiv 3 \pmod{4}$ and consider the set

$$S = \{j^2 \bmod p \mid j \in \mathbf{Z}_p \setminus \{0\}\},$$

of all quadratic residues modulo p . Note that for every nonzero element $a \in \mathbf{Z}_p$, the set $aS \pmod{p}$ is either the set of all quadratic residues or the set of all quadratic non-residues modulo p . The main result of Graham and Ringrose [GR90] asserts that for infinitely many primes p , the smallest quadratic nonresidue modulo p is at least $\Omega(\log p \log \log \log p)$ (this result holds for primes $p \equiv 3 \pmod{4}$ as well, as follows from the remark at the end of [GR90]). Since for such primes p , -1 is a quadratic nonresidue, it follows that for the above S and for any choice of $a, b \in \mathbf{Z}_p$, the set $aS + b$ (computed in \mathbf{Z}_p) avoids intervals of length at least $\Omega(\log p \log \log \log p)$. Choosing $m = c \log p \log \log \log p$ for an appropriate (small) constant c , and defining $n = \lceil p/m \rceil$, it follows that $|S| = (p - 1)/2 = \Omega(n \log n \log \log \log n)$ is not mapped onto $[0, n - 1]$ under any choice of a and b .

A final question is whether there exists a class \mathcal{H} of size only $2^{O(\log \log |U| + \log n)}$ and with $L_n^n(\mathcal{H}) = O(\log n / \log \log n)$. Note that linear maps over \mathbf{Z}_2 , even combined with collapsing the universe, use $O(\log \log |U| + (\log n)^2)$ random bits while the simple scheme using higher degree polynomials uses $O(\log \log |U| + (\log n)^2 / \log \log n)$.

References

- [ABI86] N. Alon, L. Babai and A. Itai, A fast and simple randomized parallel algorithm for the maximal independent set problem. *J.*

Algorithms **7** (1986) 567–583.

- [AS92] N. Alon and J. H. Spencer, *The Probabilistic Method*, Wiley, New York, 1992.
- [AHNR95] A. Andersson, T. Hagerup, S. Nilsson, and R. Raman, Sorting in linear time?, in: *Proc. 27th ACM Symposium on Theory of Computing*, 1995, pp. 427–436.
- [CW79] J. L. Carter and M. N. Wegman, Universal classes of hash functions, *J. Comput. Syst. Sci.* **18** (1979) 143–154.
- [CLR90] T. H. Cormen, C. E. Leiserson, and R. L. Rivest, *Introduction to Algorithms*, MIT Press, 1990.
- [DHKP93] M. Dietzfelbinger, T. Hagerup, J. Katajainen, and M. Penttonen, A reliable randomized algorithm for the closest-pair problem, Technical Report 513, Fachbereich Informatik, Universität Dortmund, 1993.
- [DM90] M. Dietzfelbinger and F. Meyer auf der Heide, Dynamic hashing in real time, in: J. Buchmann, H. Ganzinger, W. J. Paul (Eds.): *Informatik · Festschrift zum 60. Geburtstag von Günter Hotz*, Teubner-Texte zur Informatik, Band 1, B. G. Teubner, 1992, pp. 95–119. (A preliminary version appeared under the title “A New Universal Class of Hash Functions and Dynamic Hashing in Real Time” in *ICALP’90*.)
- [DKMHRT94] M. Dietzfelbinger, A. Karlin, K. Mehlhorn, F. Meyer Auf Der Heide, H. Rohnert, R.E. Tarjan, Dynamic perfect hashing: upper and lower bounds, *SIAM J. Comput.* **23** (1994) 738–761.
- [DGMP92] M. Dietzfelbinger, J. Gil, Y. Matias, and N. Pippenger, Polynomial hash functions are reliable, *ICALP’92*, Springer LNCS 623, pp. 235–246.
- [FKS84] M. L. Fredman, J. Komlós, and E. Szemerédi, Storing a sparse table with $O(1)$ worst case access time, *J. Ass. Comput. Mach.* **31** (1984) 538–544.

- [GBY90] G. Gonnet and R. Baeza-Yates, *Handbook of Algorithms and Data Structures*, Addison-Wesley, 1991.
- [GR90] S.W. Graham and C.J. Ringrose, Lower bounds for least quadratic nonresidues, in: *Analytic Number Theory: Proceedings of a Conference in Honor of P.T. Bateman*, B.C. Berndt et al. (Eds.), Birkhäuser, Boston, 1990.
- [MCW78] G. Markowsky, J.L. Carter, and M.N. Wegman, Analysis of a universal class of hash functions, in: *Proc. 7th Conference on Math. Found. of Computer Science (MFCS)*, 1978, Springer LNCS 64, pp. 345–354.
- [MV84] K. Mehlhorn and U. Vishkin, Randomized and deterministic simulations of PRAMs by parallel machines with restricted granularity of parallel memories., *Acta Informatica* **21** (1984) 339–374.
- [MNT93] Y. Mansour, N. Nisan, and P. Tiwari, The computational complexity of universal hashing. *Theoretical Computer Science* **107** (1993) 121–133.
- [PA95] J. Pach and P.K. Agarwal, *Combinatorial Geometry*, Wiley 1995.
- [S89] A. Siegel, On universal classes of fast high performance hash functions, their time-space tradeoff, and their application, in: *Proc. 30th IEEE Symposium on Foundations of Computer Science*, 1989, pp. 20–25.
- [VC71] V.A. Vapnik and A.Y. Chervonenkis, On the uniform convergence of relative frequencies of events to their probabilities, *Theory of Prob. Applications* **16** (1971) 264–280.

Recent BRICS Report Series Publications

- RS-97-16 Noga Alon, Martin Dietzfelbinger, Peter Bro Miltersen, Erez Petrank, and Gábor Tardos. *Linear Hashing*. June 1997. 22 pp. A preliminary version appeared with the title *Is Linear Hashing Good?* in *The Twenty-ninth Annual ACM Symposium on Theory of Computing, STOC '97*, pages 465–474.
- RS-97-15 Pierre-Louis Curien, Gordon Plotkin, and Glynn Winskel. *Bistructures, Bidomains and Linear Logic*. June 1997. 41 pp.
- RS-97-14 Arne Andersson, Peter Bro Miltersen, Søren Riis, and Mikkel Thorup. *Dictionaries on AC^0 RAMs: Query Time $\Theta(\sqrt{\log n / \log \log n})$ is Necessary and Sufficient*. June 1997. 18 pp. Appears in *37th Annual Symposium on Foundations of Computer Science, FOCS '96 Proceedings*, pages 441–450.
- RS-97-13 Jørgen H. Andersen and Kim G. Larsen. *Compositional Safety Logics*. June 1997. 16 pp.
- RS-97-12 Andrej Brodnik, Peter Bro Miltersen, and J. Ian Munro. *Trans-Dichotomous Algorithms without Multiplication — some Upper and Lower Bounds*. May 1997. 19 pp. Appears in Dehne, Rau-Chaulin, Sack and Tamassio, editors, *Algorithms and Data Structures: 5th International Workshop, WADS '97 Proceedings*, LNCS 1272, 1997, pages 426–439.
- RS-97-11 Kārlis Čerāns, Jens Chr. Godskesen, and Kim G. Larsen. *Timed Modal Specification — Theory and Tools*. April 1997. 32 pp.
- RS-97-10 Thomas Troels Hildebrandt and Vladimiro Sassone. *Transition Systems with Independence and Multi-Arcs*. April 1997. 20 pp. Appears in Peled, Pratt and Holzmann, editors, *DIMACS Workshop on Partial Order Methods in Verification, POMIV '96*, pages 273–288.
- RS-97-9 Jesper G. Henriksen and P. S. Thiagarajan. *A Product Version of Dynamic Linear Time Temporal Logic*. April 1997. 18 pp. Appears in Mazurkiewicz and Winkowski, editors, *Concurrency Theory: 8th International Conference, CONCUR '97 Proceedings*, LNCS 1243, 1997, pages 45–58.
- RS-97-8 Jesper G. Henriksen and P. S. Thiagarajan. *Dynamic Linear Time Temporal Logic*. April 1997. 33 pp.