

FINITE TIME BLOW UP FOR SYSTEMS OF WAVE EQUATIONS WITH MULTIPLE SPEEDS POSED ON ASYMPTOTICALLY EUCLIDEAN MANIFOLD

MENGYUN LIU

ABSTRACT. In this work, we study the finite time blow-up phenomenon of three types of semilinear wave systems with multiple speeds, posed on asymptotically Euclidean manifolds. We establish the upper bound estimates for the lifespan of solutions when the spatial dimension $n \geq 2$. In particular, for system related to the Glassey conjecture, we obtained the finite time blow up results under a new curve. This new curve is sharp at least for $n = 3$.

1. INTRODUCTION

In this paper, we study the finite time blow up phenomenon of three kinds of semilinear wave systems with multiple speeds, posed on asymptotically Euclidean (Riemannian) manifold $(\mathbb{R}^n, \mathbf{g})$ with $n \geq 2$. Before preceding, we give the definition of asymptotically Euclidean. We mean that $(\mathbb{R}^n, \mathbf{g})$ is certain perturbation of the Euclidean space $(\mathbb{R}^n, \mathbf{g}_0)$. More precisely, we assume \mathbf{g} can be decomposed as

$$(1.1) \quad \mathbf{g} = \mathbf{g}_1 + \mathbf{g}_2 ,$$

where \mathbf{g}_1 is a spherical symmetric, long range perturbation of \mathbf{g}_0 , and \mathbf{g}_2 is an exponential (short range) perturbation. By definition, there exists polar coordinates (r, ω) for $(\mathbb{R}^n, \mathbf{g}_1)$, in which we can write

$$(1.2) \quad \mathbf{g}_1 = K^2(r)dr^2 + r^2d\omega^2 ,$$

where $d\omega^2$ is the standard metric on the unit sphere \mathbb{S}^{n-1} , and

$$(1.3) \quad |\partial_r^m(K-1)| \lesssim \langle r \rangle^{-m-\rho}, m = 0, 1, 2.$$

for some given constant $\rho > 0$. Here and in what follows, $\langle x \rangle = \sqrt{1 + |x|^2}$, and we use $A \lesssim B$ ($A \gtrsim B$) to stand for $A \leq CB$ ($A \geq CB$) where the constant C may change from line to line. Equipped with the coordinates $x = r\omega$, we have

$$\mathbf{g} = g_{jk}(x)dx^j dx^k \equiv \sum_{j,k=1}^n g_{jk}(x)dx^j dx^k , \quad \mathbf{g}_2 = g_{2,jk}(x)dx^j dx^k ,$$

where we have used the convention that Latin indices j, k range from 1 to n and the Einstein summation convention for repeated upper and lower indices. Concerning

Date: November 30, 2023.

2010 Mathematics Subject Classification. 58J45, 58J05, 35L71, 35B40, 35B33, 35B44, 35B09, 35L05.

Key words and phrases. multiple speeds, blow up, lifespan estimates, asymptotically Euclidean manifolds.

The author was supported by NSFC 12101558 and NSF of Zhejiang province LQ22A010016.

\mathbf{g}_2 , we assume it is an exponential (short range) perturbation of \mathbf{g}_1 , that is, there exists $\beta > 0$ so that

$$(1.4) \quad |\nabla g_{2,jk}| + |g_{2,jk}| \lesssim e^{-\beta \int_0^r K(\tau) d\tau}, \quad |\nabla^2 g_{2,jk}| \lesssim 1.$$

By asymptotically Euclidean and Riemannian assumption, it is clear that there exists a constant $\delta_0 \in (0, 1)$ such that

$$(1.5) \quad \delta_0 |\xi|^2 \leq g^{jk} \xi_j \xi_k \leq \delta_0^{-1} |\xi|^2, \forall \xi \in \mathbb{R}^n, \quad K \in (\delta_0, 1/\delta_0).$$

The first kind of system that we studied is semilinear wave equation related to the Strauss conjecture. More precisely, we consider the following system with multiple speeds posed on asymptotically Euclidean manifolds (1.1)-(1.5).

$$(1.6) \quad \begin{cases} \partial_t^2 u - c^2 \Delta_{\mathbf{g}} u = |v|^p, (t, x) \in (0, T) \times \mathbb{R}^n, \\ \partial_t^2 v - \Delta_{\mathbf{g}} v = |u|^q, \\ u(0, x) = \varepsilon u_0(x), \quad u_t(0, x) = \varepsilon u_1(x) \\ v(0, x) = \varepsilon v_0(x), \quad v_t(0, x) = \varepsilon v_1(x) \end{cases}$$

Here $1 < p, q \in \mathbb{R}$, $0 < c \in \mathbb{R}$, $(u_0, u_1), (v_0, v_1) \in C_c^\infty(\mathbb{R}^n)$ and the small parameter $\varepsilon > 0$ measures the size of the data. As usual, to show blow up, we assume both (u_0, u_1) and (v_0, v_1) are nontrivial, nonnegative and supported in $B_R := \{x \in \mathbb{R}^n : |x| = r \leq R\}$ for some $R > 0$. It has been well investigated that the critical exponent $p_S(n)$ to Strauss problem

$$\partial_t^2 u - \Delta u = |u|^p,$$

is the positive root of the equation

$$(n-1)p^2 - (n+1)p - 2 = 0.$$

As a natural generalization, there is interest in determining whether a critical curve exists for the corresponding system (1.6).

When $c = 1$ and $\mathbf{g} = \mathbf{g}_0$, the critical curve has been well studied by Santo-Georgiev-Mitidieri [18]. They showed that the curve

$$\Gamma_{SS}(p, q, n) := \max\left\{\frac{p+2+q^{-1}}{pq-1}, \frac{q+2+p^{-1}}{pq-1}\right\} - \frac{n-1}{2} = 0, \quad n \geq 2,$$

plays the same role of $p_S(n)$ to Strauss problem at least in a neighborhood of $(p_S(n), p_S(n))$. More precisely, they obtained the finite time blow up results when $\Gamma_{SS}(p, q, n) > 0$, $n \geq 1$ and the global existence when $\Gamma_{SS}(p, q, n) < 0$, $n \geq 2$ under some additional restriction on p, q . See also [16] for case $n = 3$ and blow up result in [4]. For the critical case $\Gamma_{SS}(p, q, n) = 0$, it has also been proved there is no global existence in low space dimensions. See [17], [1], [11] [10] for $n = 3$ and [10] for $n = 2$. Recently, Ikeda-Sobojima-Wakasa [8] obtained upper bound estimates of the lifespan when $\Gamma_{SS}(p, q, n) \geq 0$, $n \geq 2$.

When the propagation speeds are different, that is $c \neq 1$, the existing results are all in low dimensional Euclidean space $\mathbf{g} = \mathbf{g}_0$. If $p \leq q$, it is easy to see that

$$\Gamma_{SS}(p, q, n) = \frac{q+2+p^{-1}}{pq-1} - \frac{n-1}{2} = 0,$$

in which case Kubo-Ohta [11], [12] proved that the critical curve is still $\Gamma_{SS}(p, q, n) = 0$ in low dimensions $n = 2, 3$. More precisely, they obtained the global existence when $\Gamma_{SS}(p, q, n) < 0$ and the finite time blow up results when $\Gamma_{SS}(p, q, n) \geq 0$. To the best of the authors' knowledge, there is no results for higher dimensions. In this

paper, we obtain the finite time blow up results with $\Gamma_{SS}(p, q, n) > 0, n \geq 2$ as well as upper bound estimates of the lifespan, to system (1.6), for any $c > 0$, posed on asymptotically Euclidean manifolds.

For the strategy of proof, we turn the systems into ODE systems by defining suitable functionals. For those ODE systems, we apply iteration methods to give upper bound estimates of the lifespan. The main innovation is the existence of a class of generalized “eigenfunctions” for the Laplace-Beltrami operator, $\Delta_{\mathbf{g}}\phi_\lambda = \lambda^2\phi_\lambda$, with small parameter $\lambda \in (0, \lambda_0)$ and desired asymptotical behavior, see Lemma 2.1. As usual, to show blow up, we need the solution to satisfy finite speed of propagation. For the linear equation $\partial_t^2 u - \Delta_{\mathbf{g}_1} u = 0$, with compacted initial data, the support of solution u satisfies

$$(1.7) \quad \text{supp } u \subset \{(t, x); \int_0^{|x|} K(\tau) d\tau \leq t + R_1\} .$$

As \mathbf{g}_2 is short-range perturbation, which does not affect speed of propagation too much, we still have (1.7) to $\partial_t^2 u - \Delta_{\mathbf{g}} u = 0$, for some $R_1 \geq \int_0^{R_0} K(\tau) d\tau$. Hence we assume the support of the solutions to system (1.6) satisfies

$$(1.8) \quad \begin{cases} \text{supp } u \subset D_1 =: \{(t, x); \int_0^{|x|} K(\tau) d\tau \leq ct + R_1\} , \\ \text{supp } v \subset D_2 =: \{(t, x); \int_0^{|x|} K(\tau) d\tau \leq t + R_1\} . \end{cases}$$

Theorem 1.1. *Let $n \geq 2$, $0 < c \in \mathbb{R}$ and $\Gamma_{SS}(p, q, n) > 0$. Consider the system (1.6) posed on asymptotically Euclidean manifolds (1.1)-(1.5). Assuming that the data are nontrivial and satisfying*

$$(1.9) \quad \begin{cases} \int u_i dv_{\mathbf{g}} > 0, \int_{\mathbb{R}^n} u_0 \phi_\lambda(x) dv_{\mathbf{g}} > 0, \int_{\mathbb{R}^n} (u_1 - c\lambda u_0) \phi_\lambda(x) dv_{\mathbf{g}} > 0, \\ \int v_i dv_{\mathbf{g}} > 0, \int_{\mathbb{R}^n} v_0 \phi_\lambda(x) dv_{\mathbf{g}} > 0, \int_{\mathbb{R}^n} (v_1 - \lambda v_0) \phi_\lambda(x) dv_{\mathbf{g}} > 0, \end{cases}$$

where $i = 0, 1$. Suppose it has a weak solution $u, v \in C^2([0, T_\varepsilon]; \mathcal{D}'(\mathbb{R}^n))$ with $|v|^p, |u|^q \in C([0, T_\varepsilon]; \mathcal{D}'(\mathbb{R}^n))$ and satisfying (1.8). Then there exist a constant $\varepsilon_0 > 0$, such that for any $\varepsilon \in (0, \varepsilon_0)$, the solution will blow up at finite time. Moreover, we have the following estimate for the lifespan T_ε

$$T_\varepsilon \lesssim \varepsilon^{-\Gamma_{SS}(p, q, n)^{-1}} .$$


Remark 1.1. Kubo-Ohta [11], [12] proved that the critical curve is $\Gamma_{SS}(p, q, n) = 0$ in low dimensions $n = 2, 3$ whenever $0 < c \in \mathbb{R}$. Based on our result, it is very natural to conjecture that for any $c > 0$ and $n \geq 2$, the critical curve to (1.6) is $\Gamma_{SS}(p, q, n) = 0$.

The second kind of system that we considered is closely related to the Glassey conjecture.

$$(1.10) \quad \begin{cases} \partial_t^2 u - c^2 \Delta_{\mathbf{g}} u = |v_t|^p, (t, x) \in (0, T) \times \mathbb{R}^n, \\ \partial_t^2 v - \Delta_{\mathbf{g}} v = |u_t|^q, \\ u(0, x) = \varepsilon u_0(x), u_t(0, x) = \varepsilon u_1(x) \\ v(0, x) = \varepsilon v_0(x), v_t(0, x) = \varepsilon v_1(x) \end{cases}$$

Similarly, we assume $(u_0, u_1), (v_0, v_1) \in C_c^\infty(\mathbb{R}^n)$ and they are nontrivial, nonnegative and supported in B_R . It well known that the critical exponent to

$$\partial_t^2 u - \Delta u = |u_t|^p ,$$



is $p_G(n) = 1 + 2/(n-1)$. It is also interesting to determine the critical curve of the corresponding system (1.10).

When $c = 1$ and $\mathbf{g} = \mathbf{g}_0$, Deng [4] has obtained the blow up results of system (1.10) under the condition

$$1 < pq < \infty, \begin{cases} 1 \leq p < \infty, n = 1, \\ 1 < p \leq 2, n = 2, \\ p = 1, n = 3, \end{cases}$$

and

$$1 < pq \leq \frac{(n+1)p}{(n-1)p-2}, \begin{cases} 2 < p \leq 3, n = 2, \\ 1 < p \leq 2, n = 3, \\ 1 < p \leq \frac{n+1}{n-1}, n \geq 4. \end{cases}$$

Recently, Ikeda-Sobojima-Wakasa [8] obtained the upper bound estimates of lifespan on a larger range (except $n = 1$)

$$(1.11) \quad \Gamma_{GG}(p, q, n) := \max\left\{\frac{p+1}{pq-1}, \frac{q+1}{pq-1}\right\} - \frac{n-1}{2} \geq 0, \quad p, q > 1, n \geq 2.$$

For global existence, Kubo-Kubota-Sunagawa [6] obtained the radially symmetric global solution in three spatial dimensions under the conditions

$$(1.12) \quad \frac{q+1}{pq-1} < 1, \quad 1 < p \leq q, \quad n = 3.$$

It can be observed that the condition $\Gamma_{GG}(p, q, n) \geq 0$ is sharp, at least for $n = 3$.

However, when the wave speed $c \neq 1$, the critical curve of (1.10) has the potential to change. Kubo-Kubota-Sunagawa [6] obtained the radially symmetric global solution when $\mathbf{g} = \mathbf{g}_0$ and

$$(1.13) \quad \frac{q}{pq-1} < 1, \quad 1 < p \leq q, \quad n = 3.$$

Compare with (1.12) (see figure 1), we are convinced that the wave speed does effect the critical curve of (1.10). For the blow up part, there is few results. Xu [19] obtained the blow up results under the condition

$$(1.14) \quad \frac{p+q+2}{pq-1} \geq 2n,$$

when $\mathbf{g} = \mathbf{g}_0$. We can see that there is a gap between (1.14) and (1.13) when $n = 3$. In this work, we fill the gap and find that the weak solution of system (1.10) under

the condition

$$(1.15) \quad \Gamma_{GG}^*(p, q, n) = \max\left\{\frac{p}{pq-1}, \frac{q}{pq-1}\right\} - \frac{n-1}{2} > 0, n \geq 2,$$

will blow up at finite time whenever $c \neq 1$.

Theorem 1.2. *Let $n \geq 2$, $0 < c \in \mathbb{R}$ and*

$$(1.16) \quad \begin{cases} \Gamma_{GG}^*(p, q, n) > 0, c \neq 1, \\ \Gamma_{GG}(p, q, n) > 0, c = 1. \end{cases}$$

Consider the system (1.10) posed on asymptotically Euclidean manifolds (1.1)-(1.5). Assuming that the data are nontrivial and satisfying

$$(1.17) \quad \begin{cases} \int_{\mathbb{R}^n} (u_1 - c\lambda u_0) \phi_\lambda(x) dv_{\mathbf{g}} > 0, \\ \int_{\mathbb{R}^n} (v_1 - \lambda v_0) \phi_\lambda(x) dv_{\mathbf{g}} > 0. \end{cases}$$

Suppose it has a weak solution $u, v \in C^2([0, T_\varepsilon]; \mathcal{D}'(\mathbb{R}^n))$ with $|v_t|^p, |u_t|^q \in C([0, T_\varepsilon]; \mathcal{D}'(\mathbb{R}^n))$ and satisfying (1.8). Then there exists a constant $\varepsilon_0 > 0$, such that for any $\varepsilon \in (0, \varepsilon_0)$, the solutions will blow up at finite time. Moreover, we have the estimate for the lifespan T_ε

$$(1.18) \quad T_\varepsilon \lesssim \begin{cases} \varepsilon^{-\Gamma_{GG}(q, p, n)^{-1}}, c = 1, \\ \varepsilon^{-\Gamma_{GG}^*(p, q, n)^{-1}}, c \neq 1. \end{cases}$$

Remark 1.2. Based on the work of Kubo-Kubota-Sunagawa [6], we conjecture that the critical curve to system (1.6) is $\Gamma_{GG}^*(p, q, n) = 0$ whenever $c \neq 1$.

The last kind of system that we studied is related to both the Strauss conjecture and the Glassey conjecture. More precisely, we consider the following system

$$(1.19) \quad \begin{cases} \partial_t^2 u - c^2 \Delta_{\mathbf{g}} u = |v|^p, (t, x) \in (0, T) \times \mathbb{R}^n, \\ \partial_t^2 v - \Delta_{\mathbf{g}} v = |u_t|^q, (t, x) \in (0, T) \times \mathbb{R}^n, \\ u(0, x) = \varepsilon u_0(x), u_t(0, x) = \varepsilon u_1(x), \\ v(0, x) = \varepsilon v_0(x), v_t(0, x) = \varepsilon v_1(x). \end{cases}$$

When $c = 1$ and $\mathbf{g} = \mathbf{g}_0$, Hidano-Yokoyama [5] obtained the finite time blow up result when

$$\frac{p+1+q^{-1}}{pq-1} - \frac{n-1}{2} > 0, q < 2n/(n-1), n \geq 2.$$

Later, Ikeda-Sobojima-Wakasa [8] obtained the finite time blow up as well as the upper bound estimates of lifespan on a larger range

$$(1.20) \quad \Gamma_{SG}(p, q, n) := \max\left\{\frac{p+1+q^{-1}}{pq-1}, \frac{2+p^{-1}}{pq-1}\right\} - \frac{n-1}{2} \geq 0, n \geq 2.$$

Dai-Fang-Wang [3] showed that there are no global weak solutions of (1.19) when

$$(1.21) \quad \frac{q+1}{pq-1} - \frac{n-1}{2} > 0, \frac{(n-1)(p-1)}{2} < 1, n \geq 2,$$

and the global existence in three dimension with

$$\frac{q+1+1/(pq)}{pq-1} < 1, 1 < p < 2, 2 < q < 3, n = 3.$$

To the best of the author's knowledge, there are no results for multiple speeds cases. In this paper, we obtained the blow up results of system (1.19) with

$$\begin{cases} M(p, q, n) := \min\{\Gamma_{GG}(p, q, n), \Gamma_{SG}(p, q, n)\} > 0, & c = 1, \\ M^*(p, q, n) := \min\{\Gamma_{GG}^*(p, q, n), \Gamma_{SG}(p, q, n)\} > 0, & c \neq 1. \end{cases}$$

Theorem 1.3. *Let $n \geq 2$, $0 < c \in \mathbb{R}$ and*

$$(1.22) \quad \begin{cases} M(p, q, n) > 0, & c = 1, \\ M^*(p, q, n) > 0, & c \neq 1. \end{cases}$$

Consider the system (1.19) posed on asymptotically Euclidean manifolds (1.1)-(1.5). Assuming that the data are nontrivial satisfying

$$(1.23) \quad \begin{cases} \int u_i dv_{\mathbf{g}} > 0, \int_{\mathbb{R}^n} u_0 \phi_{\lambda}(x) dv_{\mathbf{g}} > 0, \int_{\mathbb{R}^n} (u_1 - c\lambda u_0) \phi_{\lambda}(x) dv_{\mathbf{g}} > 0, \\ \int v_i dv_{\mathbf{g}} > 0, \int_{\mathbb{R}^n} v_0 \phi_{\lambda}(x) dv_{\mathbf{g}} > 0, \int_{\mathbb{R}^n} (v_1 - \lambda v_0) \phi_{\lambda}(x) dv_{\mathbf{g}} > 0, \end{cases}$$

where $i = 0, 1$. Suppose it has a weak solution $u, v \in C^2([0, T_{\varepsilon}); \mathcal{D}'(\mathbb{R}^n))$ with $|v|^p, |u_t|^q \in C([0, T_{\varepsilon}); \mathcal{D}'(\mathbb{R}^n))$ and satisfying (1.8). Then there exists a constant $\varepsilon_0 > 0$, such that for any $\varepsilon \in (0, \varepsilon_0)$, the solutions will blow up at finite time. Moreover, we have the estimate for the lifespan T_{ε}

$$T_{\varepsilon} \lesssim \begin{cases} \varepsilon^{-M(p, q, n)^{-1}}, & M(p, q, n) > 0, \quad c = 1, \\ \varepsilon^{-M^*(p, q, n)^{-1}}, & M^*(p, q, n) > 0, \quad c \neq 1. \end{cases}$$

Outline. Our paper is organized as follows. In Section 2, we list the existence results of special solutions for the elliptic ‘‘eigenvalue’’ problems (2.1), with certain asymptotic behavior. These solutions play important role in constructing the test functions and the proof of blow up results. Moreover, we list a Lemma in [18] for ODE system which can be viewed as a vector version of Kato type Lemma. In Sections 3-5, we give the proof of Theorems 1.1-1.3 by applying iteration method (see, e.g., [13]).

2. PRELIMINARY

In this section, we list some Lemmas we shall use later. We first construct special positive standing wave solutions, of the form

$$\psi_1(t, x) = e^{-c\lambda t} \phi_{\lambda}(x), \quad \psi_2(t, x) = e^{-\lambda t} \phi_{\lambda}(x),$$

to the linear problem,

$$\partial_t^2 u - c^2 \Delta_{\mathbf{g}} u = 0, \quad \partial_t^2 v - \Delta_{\mathbf{g}} v = 0,$$

with the desired asymptotic behavior. In turn, it is reduced to constructing solutions to certain elliptic ‘‘eigenvalue’’ problems:

$$(2.1) \quad \Delta_{\mathbf{g}} \phi_{\lambda} = \lambda^2 \phi_{\lambda}.$$

These solutions will play a key role in the construction of the test functions.

Lemma 2.1 (Lemma 3.1 in [14]). *Let $n \geq 2$ and $(\mathbb{R}^n, \mathbf{g})$ be asymptotically Euclidean manifold with (1.1)-(1.5). Then there exist $\lambda_0, c_0 > 0$ such that for any $0 < \lambda \leq \lambda_0$, there is a solution of (2.1) satisfying*

$$(2.2) \quad c_0 < \phi_{\lambda}(x) < c_0^{-1} \langle \lambda r \rangle^{-\frac{n-1}{2}} e^{\lambda \int_0^r K(\tau) d\tau}.$$

With the asymptotic behavior of ϕ_λ in hand, we could have the following estimates for ψ_1, ψ_2 .

Lemma 2.2. *Let $n \geq 2$, $p, q > 1$. Then we have*

$$\begin{aligned} \int_{D_i} \psi_i^p dv_{\mathbf{g}} &\lesssim (t+1)^{n-1-\frac{n-1}{2}p}, i = 1, 2, \\ \int_{D_2} \psi_1^{-\frac{p'}{p}} \psi_2^{p'} dv_{\mathbf{g}} &\lesssim e^{\frac{\lambda(c-1)t}{p-1}} (t+1)^{\frac{n-1}{2}}, \\ \int_{D_1} \psi_2^{-\frac{q'}{q}} \psi_1^{q'} dv_{\mathbf{g}} &\lesssim e^{\frac{\lambda(1-c)t}{q-1}} (t+1)^{\frac{n-1}{2}}. \end{aligned}$$

Proof. We only give a short proof of the second inequality since others follow the same way.

$$\int_{D_2} \psi_1^{-\frac{p'}{p}} \psi_2^{p'} dv_{\mathbf{g}} = e^{\frac{\lambda t(c-p)}{p-1}} \int_{D_2} \phi_\lambda(x) dv_{\mathbf{g}} = e^{\frac{\lambda(c-1)t}{p-1}} \int_{D_2} e^{-\lambda t} \phi_\lambda dv_{\mathbf{g}}.$$

Then by Lemma 3.2 in [15] (with $q = 1$), we have that

$$\int_{D_2} e^{-\lambda t} \phi_\lambda dv_{\mathbf{g}} \lesssim (t+1)^{\frac{n-1}{2}},$$

which complete the proof. For the reader's convenience, we present a proof here. We divide the region D_2 into two disjoint parts: $D_2 = N_1 \cup N_2$ where

$$N_1 = \{(t, x); \int_0^{|x|} K(\tau) d\tau \leq \frac{t+R_1}{2}\}.$$

For the region N_1 , we have

$$\int_{N_1} \phi_\lambda dv_{\mathbf{g}} \lesssim e^{-\lambda t} \int_{N_1} (1 + \lambda|x|)^{-\frac{n-1}{2}} e^{\lambda \int_0^{|x|} K(\tau) d\tau} dv_{\mathbf{g}}.$$

Let $\tilde{r} = \int_0^{|x|} K(\tau) d\tau$, then $d\tilde{r} = K(r)dr$ and $\delta_0 r \leq \tilde{r} \leq r/\delta_0$ since $K \in [\delta_0, 1/\delta_0]$. Then we get

$$\begin{aligned} \int_{N_1} \phi_\lambda dv_{\mathbf{g}} &\lesssim e^{-\lambda t} \int_0^{\frac{t+R_1}{2}} (1 + \tilde{r})^{n-1-\frac{n-1}{2}} e^{\lambda \tilde{r}} d\tilde{r} \\ &\lesssim e^{-\lambda t} \int_0^{\frac{t+R_1}{2}} e^{\frac{3}{2}\lambda \tilde{r}} d\tilde{r} \\ &\lesssim e^{-\frac{\lambda}{4}t} \lesssim (1+t)^{\frac{n-1}{2}}, \end{aligned}$$

where we have used the fact that e^{-t} decays faster than any polynomial. For the region N_2 , it is easy to see

$$\begin{aligned} \int_{N_2} \phi_\lambda dv_{\mathbf{g}} &\lesssim e^{-\lambda t} \int_{\frac{t+R_1}{2}}^{t+R_1} e^{\lambda \tilde{r}} (1 + \tilde{r})^{n-1-\frac{n-1}{2}} d\tilde{r} \\ &\lesssim (1+t)^{\frac{n-1}{2}} \int_{\frac{t+R_1}{2}}^{t+R_1} e^{\lambda \tilde{r} - \lambda t} d\tilde{r} \\ &\lesssim (1+t)^{\frac{n-1}{2}}. \end{aligned}$$

□

At the end of this section, we list a Lemma for ODE systems to blow up in finite time. This Lemma can be viewed as a vector version of the Kato type Lemma.

Lemma 2.3 (Lemma 2.1 in [18]). *Let $a, b \in [0, \infty]$ with $a < b$ and $M(t), N(t) \in C([a, b]; \mathbb{R})$. Suppose that for all $t \in [a, b]$, we have*

$$(2.3) \quad \begin{cases} M(t) \geq C(t+R) , \\ N(t) \geq C(t+R)^s , \\ M''(t) \geq C(t+R)^{-\alpha}(N(t))^e , \\ N''(t) \geq C(t+R)^{-\beta}(M(t))^l , \end{cases}$$

where $C, R > 0$, $s \geq 1$, $\alpha, \beta > 0$, $e, l > 0$. Moreover, if $el > 1$ and

$$l(\alpha - 2) + \beta - 2 < s(el - 1) ,$$

then $b < \infty$.

3. PROOF OF THEOREMS 1.1

In this section, we give the proof the Theorem 1.1. Without loss of generality, we assume $0 < c \leq 1$. In this case, we see that $D_1 \subset D_2$. We first define functionals

$$F(t) = \int_{D_2} u dv_{\mathbf{g}} , \quad G(t) = \int_{D_2} v dv_{\mathbf{g}} .$$

Then by the system (1.6), we know that

$$F''(t)^1 = \int_{D_2} |v|^p dv_{\mathbf{g}} , \quad G''(t) = \int_{D_2} |u|^q dv_{\mathbf{g}} .$$

To connect F'' , G'' with F , G , we apply Hölder's inequality to get

$$F'' = \int_{D_2} |v|^p dv_{\mathbf{g}} \gtrsim \frac{(\int_{D_2} v dv_{\mathbf{g}})^p}{(t+1)^{n(p-1)}} , \quad G'' = \int_{D_2} |u|^q dv_{\mathbf{g}} \gtrsim \frac{(\int_{D_2} u dv_{\mathbf{g}})^q}{(t+1)^{n(q-1)}}$$

Hence we can get the following ODE system

$$(3.1) \quad F'' \geq c_0 \frac{G^p}{(t+1)^{n(p-1)}} , \quad G'' \geq c_1 \frac{F^q}{(t+1)^{n(q-1)}} ,$$

for some constants $c_0, c_1 > 0$. By the assumption (1.9) on the initial data, we have the lower bound of F , G

$$(3.2) \quad F(t) \gtrsim \varepsilon(t+1) , \quad G \gtrsim \varepsilon(t+1), \quad t > 0 .$$

Define auxiliary functions

$$H_1(t) = \int_{D_1} u \psi_1 dv_{\mathbf{g}} , \quad H_2(t) = \int_{D_2} v \psi_2 dv_{\mathbf{g}} .$$

We will see that the lower bound of F , G could be improved to

$$(3.3) \quad F(t) \gtrsim \varepsilon^p (t+1)^{n+1-\frac{n-1}{2}p} , \quad G(t) \gtrsim \varepsilon^q (t+1)^{n+1-\frac{n-1}{2}q} , \quad t > 0 ,$$

under the condition

$$n - \frac{n-1}{2}p > 0 , \quad n - \frac{n-1}{2}q > 0 .$$

¹Note that $u|_{\partial D_1} = 0$, hence $u|_{\partial D_2} = 0$.

In fact, by (1.6), we have

$$H_2'' + 2\lambda H_2' = \int_{D_2} |u|^q \psi_2 dv_{\mathfrak{g}} \geq 0 ,$$

thus by the assumption (1.9) on the initial data ,

$$H_2'(t) \geq e^{-2\lambda t} H_2'(0) \geq 0 , H_2(t) \geq H_2(0) \gtrsim \varepsilon .$$

By applying Hölder's inequality and Lemma 2.2, we get

$$\begin{aligned} \varepsilon &\lesssim H_2 \lesssim \left(\int_{D_2} |v|^p dv_{\mathfrak{g}} \right)^{1/p} \left(\int_{D_2} |\psi_2|^{p'} dv_{\mathfrak{g}} \right)^{1/p'} \\ &\lesssim \left(\int_{D_2} |v|^p dv_{\mathfrak{g}} \right)^{1/p} \left((t+1)^{n-1-\frac{n-1}{2}p'} \right)^{1/p'} . \end{aligned}$$

Then

$$F'' = \int_{D_2} |v|^p dv_{\mathfrak{g}} \gtrsim \varepsilon^p (t+1)^{n-1-\frac{n-1}{2}p} .$$

Similarly, we see that

$$H_1'' + 2\lambda c H_1' = \int_{D_1} |v|^q \psi_1 dv_{\mathfrak{g}} \geq 0 ,$$

thus

$$H_1'(t) \geq e^{-2\lambda t} H_1'(0) \geq 0 , H_1(t) \geq H_1(0) \gtrsim \varepsilon .$$

By applying Hölder's inequality and Lemma 2.2, we get

$$\begin{aligned} \varepsilon &\lesssim H_1 \lesssim \left(\int_{D_1} |u|^q dv_{\mathfrak{g}} \right)^{1/q} \left(\int_{D_1} |\psi_1|^{q'} dv_{\mathfrak{g}} \right)^{1/q'} \\ &\lesssim \left(\int_{D_1} |u|^q dv_{\mathfrak{g}} \right)^{1/q} \left((t+1)^{n-1-\frac{n-1}{2}q'} \right)^{1/q'} , \end{aligned}$$

which yields

$$G'' = \int_{D_2} |u|^q dv_{\mathfrak{g}} \geq \int_{D_1} |u|^q dv_{\mathfrak{g}} \gtrsim \varepsilon^q (t+1)^{n-1-\frac{n-1}{2}q} .$$

Hence, we could integrate the above inequalities twice to get (3.3).

Applying Lemma 2.3 to (3.1) and (3.3), we know that the solutions will blow up if $\Gamma_{SS}(p, q, n) > 0$. In fact, by the second inequality of (3.1), we have

$$G(t) \gtrsim (t+1)^{\left(n+1-\frac{n-1}{2}p\right)q-n(q-1)+2} .$$

Heuristically, the lower bound of F could be improved by the first inequality of (3.1) if

$$\left(\left(n+1 - \frac{n-1}{2}p \right) q - n(q-1) + 2 \right) p - n(p-1) + 2 > n+1 - \frac{n-1}{2}p ,$$

similarly, the lower bound of G could be improved if

$$\left(\left(n+1 - \frac{n-1}{2}q \right) p - n(p-1) + 2 \right) q - n(q-1) + 2 > n+1 - \frac{n-1}{2}q ,$$

that is $\Gamma_{SS}(p, q, n) > 0$. Without loss of generality, we assume $p \leq q$ and in this case

$$(3.4) \quad \Gamma_{SS}(p, q, n) = \frac{p^{-1} + q + 2}{pq - 1} - \frac{n-1}{2} > 0 .$$

To get the upper bound lifespan estimates, we apply iteration method and we will see that there exists $\varepsilon_0 > 0$ such that for any $\varepsilon \in (0, \varepsilon_0)$, we have the lower bound of F, G by some sequences

$$F \geq A_j(t+1)^{a_j}, G \geq B_j(t+1)^{b_j}, j \in \mathbb{N},$$

with initial lower bound

$$A_1 = \varepsilon^p, a_1 = n+1 - \frac{n-1}{2}p.$$

In fact, by putting the lower bound of F to the second inequality of (3.1), we have

$$G'' \geq c_1 A_1^q (t+1)^{qa_1-n(q-1)} = c_1 \varepsilon^{pq} (t+1)^{qa_1-n(q-1)}.$$

Note that $G(0), G'(0) \sim \varepsilon$, then we have

$$\begin{aligned} G' &\geq \frac{c_1 \varepsilon^{pq}}{qa_1 - n(q-1) + 1} (t+1)^{qa_1-n(q-1)+1} - \frac{c_1 \varepsilon^{pq}}{qa_1 - n(q-1) + 1} + G'(0) \\ &\geq \frac{c_1 \varepsilon^{pq}}{qa_1 - n(q-1) + 1} (t+1)^{qa_1-n(q-1)+1}, \end{aligned}$$

and

$$G \geq \frac{c_1 \varepsilon^{pq} (t+1)^{qa_1-n(q-1)+2}}{(qa_1 - n(q-1) + 1)(qa_1 - n(q-1) + 2)}$$

if we take ε small enough. Hence we obtain that

$$B_1 = \frac{c_1 \varepsilon^{pq}}{b_1(b_1-1)}, b_1 = qa_1 - n(q-1) + 2,$$

and then we use above to get A_2, a_2 by the first inequality of (3.1). Following this iteration way, we can obtain the following iteration equation

$$(3.5) \quad \begin{cases} A_j = \frac{c_0 B_{j-1}^p}{(a_j-1)a_j}, a_j = pb_{j-1} - n(p-1) + 2, j \geq 2 \\ B_j = \frac{c_1 A_j^q}{(b_j-1)b_j}, b_j = qa_j - n(q-1) + 2, j \geq 1 \end{cases}$$

if we have

$$(3.6) \quad (a_j+1)A_j \leq \varepsilon, (b_j+1)B_j \leq \varepsilon, \forall j \in \mathbb{N}.$$

In fact, by (3.5) we get that

$$b_j = \left(\frac{p^{-1} + q + 2}{pq-1} - \frac{n-1}{2} \right) (pq)^j + n - \frac{2q+2}{pq-1}.$$

Since b_j and a_j are increasing to ∞ as $j \rightarrow \infty$, there exists $N_0 > 0$ such that $a_j, b_j \geq 2$ for any $j \geq N_0$. Thus we have

$$(3.7) \quad \begin{cases} (b_j+1)B_j = \frac{c_1 A_j^q}{(b_j-1)b_j} + \frac{c_1 A_j^q}{(b_j-1)} \leq 2c_1 A_j^q \leq 2c_1 c_0^q B_{j-1}^{pq}, j \geq N_0+1, \\ (a_j+1)A_j = \frac{c_0 B_{j-1}^p}{(a_j-1)a_j} + \frac{c_0 B_{j-1}^p}{(a_j-1)} \leq 2c_0 B_{j-1}^p \leq 2c_0 c_1^p A_{j-1}^{pq}, j \geq N_0+1. \end{cases}$$

For $j \leq N_0$, we take ε_0 small enough such that for any $\varepsilon \in (0, \varepsilon_0)$, we have

$$(3.8) \quad (b_j+1)B_j + (a_j+1)A_j \leq \frac{\varepsilon}{2c_0 c_1^p + 2c_1 c_0^q + 1}.$$

Combine (3.8) and (3.7), we get (3.6).

For the coefficient B_j , it is easy to see that

$$B_j = \frac{c_1 c_0^q B_{j-1}^{pq}}{((a_j - 1)a_j)^q (b_j - 1)b_j} \geq \frac{c_1 c_0^q B_{j-1}^{pq}}{(pb_j + 2)^{2q+2}} \geq \frac{M B_{j-1}^{pq}}{(pq)^{j(2q+2)}} ,$$

where

$$M = c_1 c_0^q \left(2 + \frac{n+1}{2}p + \frac{1+pq+2p}{pq-1} \right)^{-(2q+2)} .$$

Hence we get

$$\begin{aligned} \ln B_j &\geq (pq) \ln B_{j-1} + \ln M - j(2q+2) \ln(pq) \\ &\geq (pq)^2 \ln B_{j-2} + (pq+1) \ln M - (pq(j-1) + j)(2q+2) \ln(pq) \\ &\geq (pq)^{j-1} \ln B_1 - (pq)^{j-1} \sum_{k=1}^{j-1} \frac{(k+1)(2q+2) \ln(pq) - \ln M}{(pq)^k} . \end{aligned}$$

Note that the series is convergence, that is

$$S(\infty) = \sum_{k=1}^{\infty} \frac{(k+1)(2q+2) \ln(pq) - \ln M}{(pq)^k} < \infty .$$

Then we have

$$\ln B_j \geq (pq)^{j-1} (\ln B_1 - S(\infty)) .$$

Thus

$$\begin{aligned} G &\geq B_j(t+1)^{b_j} \\ &\geq e^{(pq)^{j-1} \left(\ln B_1 - S(\infty) + \left(\frac{2q+2}{pq-1} + q+2 - \frac{n-1}{2}pq \right) \ln(t+1) \right)} e^{(n - \frac{2q+2}{pq-1}) \ln(t+1)} , \forall j \in \mathbb{N} , \end{aligned}$$

which yields the lifespan estimate

$$T_\varepsilon \lesssim \varepsilon^{-\left(\frac{q+2+p^{-1}}{pq-1} - \frac{n-1}{2} \right)^{-1}} .$$

Similarly, if we assume $p \geq q$ that is

$$\Gamma_{SS}(p, q, n) = \frac{q^{-1} + p + 2}{pq - 1} - \frac{n - 1}{2} > 0 ,$$

and start the iteration with $B_1 = \varepsilon^q$, $b_1 = n + 1 - \frac{n-1}{2}q$, we will have that

$$T_\varepsilon \lesssim \varepsilon^{-\left(\frac{p+2+q^{-1}}{pq-1} - \frac{n-1}{2} \right)^{-1}} .$$

Hence, we obtain the lifespan estimate

$$T_\varepsilon \lesssim \varepsilon^{-\Gamma_{SS}(p, q, n)^{-1}} .$$

4. PROOF OF THEOREM 1.2

Without loss of generality, we assume $0 < c \leq 1$. In this case, we see that $D_1 \subset D_2$. We first define functionals

$$(4.1) \quad F(t) = \int_{D_2} (u_t \psi_1 + c \lambda u \psi_1) dv_{\mathbf{g}} , G(t) = \int_{D_2} (v_t \psi_2 + \lambda v \psi_2) dv_{\mathbf{g}} .$$

Then by the system (1.10), we have

$$(4.2) \quad F'(t) = \int_{D_2} |v_t|^p \psi_1 dv_{\mathbf{g}} , G'(t) = \int_{D_2} |u_t|^q \psi_2 dv_{\mathbf{g}} .$$

In order to relate F' with G , we let $H_1(t) = \int_{D_2} u_t \psi_1 dv_{\mathfrak{g}}$ and it is easy to check that

$$(2H_1 - F)' + 2c\lambda(2H_1 - F) = \int_{D_2} |v_t|^p \psi_1 dv_{\mathfrak{g}} \geq 0 ,$$

thus by the assumption on the initial data (1.17), we have

$$(2H_1 - F)(t) \geq (2H_1 - F)(0) \geq 0 ,$$

which yields

$$\frac{F}{2} \leq H_1(t) = \int_{D_2} u_t \psi_1 dv_{\mathfrak{g}} .$$

Similarly, let $H_2 = \int_{D_2} v_t \psi_2 dv_{\mathfrak{g}}$, we can get

$$\frac{G}{2} \leq H_2(t) = \int_{D_2} v_t \psi_2 dv_{\mathfrak{g}} .$$

Recall that the support of u is on D_1 by (1.8), so $H_1(t) = \int_{D_1} u_t \psi_1 dv_{\mathfrak{g}}$. By applying Hölder's inequality to H_1 and H_2 , we have

$$\begin{aligned} \int_{D_2} |v_t|^p \psi_1 dv_{\mathfrak{g}} &\gtrsim \frac{\left(\int_{D_2} v_t \psi_2 dv_{\mathfrak{g}} \right)^p}{\left(\int_{D_2} \psi_1^{-\frac{p'}{p}} \psi_2^{p'} dv_{\mathfrak{g}} \right)^{p/p'}} \gtrsim \frac{H_2^p}{\left(\int_{D_2} \psi_1^{-\frac{p'}{p}} \psi_2^{p'} dv_{\mathfrak{g}} \right)^{p/p'}} , \\ \int_{D_1} |u_t|^q \psi_2 dv_{\mathfrak{g}} &\gtrsim \frac{\left(\int_{D_1} u_t \psi_1 dv_{\mathfrak{g}} \right)^q}{\left(\int_{D_1} \psi_2^{-\frac{q'}{q}} \psi_1^{q'} dv_{\mathfrak{g}} \right)^{q/q'}} \gtrsim \frac{H_1^q}{\left(\int_{D_1} \psi_2^{-\frac{q'}{q}} \psi_1^{q'} dv_{\mathfrak{g}} \right)^{q/q'}} . \end{aligned}$$

4.1. $c = 1$. When $c = 1$, we have $D = D_1 = D_2$, by Lemma 2.2,

$$\begin{aligned} \left(\int_{D_2} \psi_1^{-\frac{p'}{p}} \psi_2^{p'} dv_{\mathfrak{g}} \right)^{p/p'} &\gtrsim (t+1)^{(n-1)(p-1)/2} , \\ \left(\int_{D_2} \psi_2^{-\frac{q'}{q}} \psi_1^{q'} dv_{\mathfrak{g}} \right)^{q/q'} &\gtrsim (t+1)^{(n-1)(q-1)/2} . \end{aligned}$$

Thus we get the following ODE system

$$(4.3) \quad \begin{cases} F'(t) \geq \frac{c_0 |G|^p}{(t+1)^{\frac{(n-1)(p-1)}{2}}} , F(0) \geq c_0 \varepsilon \\ G'(t) \geq \frac{c_1 |F|^q}{(t+1)^{\frac{(n-1)(q-1)}{2}}} , G(0) \geq c_1 \varepsilon . \end{cases}$$

where constants c_0 and c_1 depends on p, q, c, λ, n . We will see that (4.3) blows up in $\Gamma(p, q, n) > 0$. In fact, it is easy to see $F(t) \geq F(0) \geq c_0 \varepsilon$ due to $F' \geq 0$. By putting it into the second inequality in (4.3), we have

$$(4.4) \quad G'(t) \geq c_1 (c_0 \varepsilon)^q (t+1)^{-\frac{(n-1)(q-1)}{2}} ,$$

thus $G(t) \gtrsim (t+1)^{1-(n-1)(q-1)/2}$ and by (4.3) again, we get that

$$F'(t) \gtrsim (t+1)^{\left(1 - \frac{(n-1)(q-1)}{2}\right)p - \frac{(n-1)(p-1)}{2}} .$$

Heuristically, the lower bound of $F(t)$ could be improved if

$$\left(1 - \frac{(n-1)(q-1)}{2}\right)p - \frac{(n-1)(p-1)}{2} + 1 > 0 ,$$

similarly, the lower bound of $G(t)$ could be improved if

$$\left(1 - \frac{(n-1)(p-1)}{2}\right)q - \frac{(n-1)(q-1)}{2} + 1 > 0 ,$$

that is the range $\Gamma_{GG}(p, q, n) > 0$, which suggests the blow up result.

To obtain the lifespan estimates, we explore iteration argument and assume

$$(4.5) \quad F(t) \geq A_j(t+1)^{a_j} , G(t) \geq B_j(t+1)^{b_j} , j \in \mathbb{N} .$$

Without loss of generality, we assume $\frac{(n-1)(q-1)}{2} < 1$, then by (4.3), we get that

$$\begin{aligned} G(t) &\geq \frac{c_0(c_1\varepsilon)^q}{1 - \frac{(n-1)(q-1)}{2}}(t+1)^{1 - \frac{(n-1)(q-1)}{2}} - \frac{c_0(c_1\varepsilon)^q}{1 - \frac{(n-1)(q-1)}{2}} + c_1\varepsilon \\ &\geq \frac{c_0(c_1\varepsilon)^q}{1 - \frac{(n-1)(q-1)}{2}}(t+1)^{1 - \frac{(n-1)(q-1)}{2}} , t > 0 , \end{aligned}$$

if we take ε small enough. We start the iteration with

$$B_1 = \frac{c_0(c_1\varepsilon)^q}{1 - \frac{(n-1)(q-1)}{2}} , b_1 = 1 - \frac{(n-1)(q-1)}{2} ,$$

and assume

$$\Gamma_{GG}(p, q, n) = \frac{p+1}{pq-1} - \frac{n-1}{2} > 0 .$$

By putting it to (4.3), we have

$$F'(t) \geq c_0 B_1^p(t+1)^{pb_1 - \frac{(n-1)(p-1)}{2}} ,$$

that is

$$F(t) \geq \frac{c_0 B_1^p}{1 + pb_1 - \frac{(n-1)(p-1)}{2}}(t+1)^{1 + pb_1 - \frac{(n-1)(p-1)}{2}} , t > 0 ,$$

if we take ε small enough. Then we have

$$A_1 = \frac{c_0 B_1^p}{1 + pb_1 - \frac{(n-1)(p-1)}{2}} , a_1 = 1 + pb_1 - \frac{(n-1)(p-1)}{2} .$$

Following this way, we have the iteration equation

$$(4.6) \quad \begin{cases} A_j = \frac{B_j^p}{1 + pb_j - \frac{(n-1)(p-1)}{2}} , a_j = 1 + pb_j - \frac{(n-1)(p-1)}{2} , j \geq 1 , \\ B_{j+1} = \frac{A_j^q}{1 + qa_j - \frac{(n-1)(q-1)}{2}} , b_{j+1} = 1 + qa_j - \frac{(n-1)(q-1)}{2} , j \geq 1 , \end{cases}$$

If we have

$$(4.7) \quad A_j \leq c_0\varepsilon , B_j \leq c_1\varepsilon , \forall j \in \mathbb{N} .$$

In fact, by (4.6) we have

$$a_j = \left(\frac{p+1}{pq-1} - \frac{n-1}{2}\right)((pq)^j - 1) , j \in \mathbb{N} ,$$

and $a_j \rightarrow \infty$ as $j \rightarrow \infty$. Thus there exist a $N_0 > 0$, such that for any $j \geq N_0 + 1$,

$$A_j = \frac{A_{j-1}^{pq}}{a_j b_j^p} \leq A_{j-1}^{pq} , B_j = \frac{B_{j-1}^{pq}}{a_{j-1}^q b_j} \leq B_{j-1}^{pq} .$$

For $j \leq N_0$, we take $\varepsilon_0 \ll 1$ small enough to get

$$A_j \leq c_0 \varepsilon, \quad B_j \leq c_1 \varepsilon, \quad j \leq N_0,$$

for any $\varepsilon \in (0, \varepsilon_0)$.

For the sequence A_j , by definition we have that

$$A_j = \frac{A_{j-1}^{pq}}{a_j b_j^p} \geq \frac{A_{j-1}^{pq}}{K(pq)^{j(p+1)}}, \quad K = \left(q \left(\frac{p+1}{pq-1} - \frac{n-1}{2} \right) + 1 \right)^{p+1}.$$

Hence we get

$$\begin{aligned} \ln A_j &\geq (pq) \ln A_{j-1} - \ln K - j(p+1) \ln(pq) \\ &\geq (pq)^2 \ln A_{j-2} - (pq+1) \ln K - (pq(j-1) + j)(p+1) \ln(pq) \\ &\geq (pq)^{j-1} \ln A_1 - (pq)^{j-1} \sum_{k=1}^{j-1} \frac{\ln K + (k+1)(p+1) \ln(pq)}{(pq)^k}. \end{aligned}$$

Note that the series is convergence, that is

$$S(\infty) = \sum_{k=0}^{\infty} \frac{\ln K + (k+1)(p+1) \ln(pq)}{(pq)^{k+1}} < \infty.$$

Then we have

$$\ln A_j \geq (pq)^{j-1} (\ln A_1 - S(\infty)).$$

Thus

$$\begin{aligned} F &\geq A_j (t+1)^{a_j} \\ &\geq e^{(pq)^{j-1} \left(\ln A_1 - S(\infty) + \left(\frac{p+1}{pq-1} - \frac{n-1}{2} \right) pq \ln(t+1) \right)} e^{-\left(\frac{p+1}{pq-1} - \frac{n-1}{2} \right) \ln(t+1)}, \end{aligned}$$

we can see that

$$\ln A_1 - S(\infty) + \left(\frac{p+1}{pq-1} - \frac{n-1}{2} \right) pq \ln(t+1) \leq 0,$$

otherwise, $F(t) \rightarrow \infty$ as $j \rightarrow \infty$. From which we have the estimate of T_ε

$$T_\varepsilon \lesssim \varepsilon^{-\left(\frac{p+1}{pq-1} - \frac{n-1}{2} \right)^{-1}}.$$

Moreover, when $\Gamma_{GG}(p, q, n) = \frac{q+1}{pq-1} - \frac{n-1}{2} > 0$, we use the following iteration equation

$$\begin{cases} A_{j+1} = \frac{B_j^p}{a_{j+1}}, & a_{j+1} = 1 + pb_j - \frac{(n-1)(p-1)}{2}, \quad j \geq 1, \\ B_j = \frac{A_j^q}{b_j}, & b_j = 1 + qa_j - \frac{(n-1)(q-1)}{2}, \quad j \geq 1, \end{cases}$$

with

$$A_1 = \frac{c_0^p c_1 \varepsilon}{1 - \frac{(n-1)(p-1)}{2}}, \quad a_1 = 1 - \frac{(n-1)(p-1)}{2}.$$

And we can obtain another lifespan estimate

$$T_\varepsilon \lesssim \varepsilon^{-\left(\frac{q+1}{pq-1} - \frac{n-1}{2} \right)^{-1}}.$$

In summary, we have

$$(4.8) \quad T_\varepsilon \lesssim \min \left\{ \varepsilon^{-\left(\frac{p+1}{pq-1} - \frac{n-1}{2} \right)^{-1}}, \varepsilon^{-\left(\frac{q+1}{pq-1} - \frac{n-1}{2} \right)^{-1}} \right\}.$$

4.2. $0 < c < 1$. By Lemma 2.2, we have

$$(4.9) \quad \begin{cases} \left(\int_{D_2} \psi_1^{-\frac{p'}{p}} \psi_2^{p'} dv_{\mathfrak{g}} \right)^{p/p'} \lesssim e^{\lambda(c-1)t} (t+1)^{(n-1)(p-1)/2}, \\ \left(\int_{D_1} \psi_2^{-\frac{q'}{q}} \psi_1^{q'} dv_{\mathfrak{g}} \right)^{q/q'} \lesssim e^{\lambda(1-c)t} (t+1)^{(n-1)(q-1)/2}. \end{cases}$$

Thus we get the following ODE system

$$(4.10) \quad \begin{cases} F'(t) \geq \frac{c_0 |G|^p}{e^{\lambda(c-1)t} (t+1)^{\frac{(n-1)(p-1)}{2}}}, \quad F(0) \geq c_0 \varepsilon \\ G'(t) \geq \frac{c_1 |F|^q}{e^{\lambda(1-c)t} (t+1)^{\frac{(n-1)(q-1)}{2}}}, \quad G(0) \geq c_1 \varepsilon \end{cases}$$

We will see that the system (4.10) blows up in finite time when $\tilde{\Gamma}(p, q, n) > 0$. For that purpose, we set $G = H(t)e^{\lambda(c-1)t/p}$, then we have

$$(4.11) \quad F'(t) \geq \frac{c_0 |H|^p}{(t+1)^{\frac{(n-1)(p-1)}{2}}}, \quad H' + \frac{\lambda(c-1)}{p} H \geq \frac{c_1 |F|^q}{(t+1)^{\frac{(n-1)(q-1)}{2}}} e^{\lambda(c-1)t/p'}.$$

It is easy to see that $H(t) \geq H(0) = G(0) \geq c_1 \varepsilon$. By putting it into the first inequality in (4.11), we have

$$F(t) \gtrsim (t+1)^{1-(n-1)(p-1)/2},$$

then by (4.11) again, we get that

$$\begin{aligned} & e^{\lambda(c-1)t/p} H \\ & \gtrsim \int_{t/p}^t e^{\lambda(c-1)\tau} (\tau+1)^{\left(1-\frac{(n-1)(p-1)}{2}\right)q - \frac{(n-1)(q-1)}{2}} d\tau, \\ & \gtrsim (t+1)^{\left(1-\frac{(n-1)(p-1)}{2}\right)q - \frac{(n-1)(q-1)}{2}} \int_{t/p}^t e^{\lambda(c-1)\tau} d\tau \\ & \gtrsim (t+1)^{\left(1-\frac{(n-1)(p-1)}{2}\right)q - \frac{(n-1)(q-1)}{2}} e^{\lambda(c-1)t/p} \left(1 - e^{\lambda(c-1)t/p'}\right), \end{aligned}$$

Heuristically, the lower bound of $H(t)$ could be improved if

$$\left(1 - \frac{(n-1)(p-1)}{2}\right)q - \frac{(n-1)(q-1)}{2} > 0,$$

that is

$$\frac{q}{pq-1} > \frac{n-1}{2}.$$

While for the case $\frac{p}{pq-1} > \frac{n-1}{2}$, we shall give another intuition. Let $H(t) = e^{\lambda(c-1)t} F(t)$, then system (4.10) becomes

$$(4.12) \quad \begin{cases} H' + \lambda(1-c)H \geq \frac{c_0 |G|^p}{(t+1)^{(n-1)(p-1)/2}}, \quad H(0) \gtrsim \varepsilon \\ G'(t) \geq \frac{H^q}{(t+1)^{(n-1)(q-1)/2}}, \quad G(0) \geq c_1 \varepsilon \end{cases}$$

By putting $H \gtrsim \varepsilon$ into the second inequality in (4.12), we have

$$G' \gtrsim \varepsilon^q (t+1)^{-\frac{(n-1)(q-1)}{2}},$$

thus

$$G \gtrsim \varepsilon^q (t+1)^{1-\frac{(n-1)(q-1)}{2}}.$$

And then by the first inequality in (4.12), we get that

$$(e^{\lambda(1-c)t}H)' \gtrsim \varepsilon^{pq}e^{\lambda(1-c)t}(t+1)^{\left(1-\frac{(n-1)(q-1)}{2}\right)p-\frac{(n-1)(p-1)}{2}},$$

thus

$$\begin{aligned} H(t) &\gtrsim \varepsilon^{pq}(t+1)^{\left(1-\frac{(n-1)(q-1)}{2}\right)p-\frac{(n-1)(p-1)}{2}} e^{\lambda(c-1)t} \int_{t/2}^t e^{\lambda(1-c)\tau} d\tau, \\ &\gtrsim \varepsilon^{pq}(t+1)^{\left(1-\frac{(n-1)(q-1)}{2}\right)p-\frac{(n-1)(p-1)}{2}}, \end{aligned}$$

so the lower bound of H could be boosted when

$$\left(1 - \frac{(n-1)(q-1)}{2}\right)p - \frac{(n-1)(p-1)}{2} > 0,$$

which is actually

$$\frac{p}{pq-1} > \frac{n-1}{2}.$$

To get the upper bound of lifespan estimates, we explore iteration argument. Without loss of generality, we assume $p \geq q$ and in this case

$$\Gamma_{GG}^*(p, q, n) = \frac{p}{pq-1} > 0.$$

We assume

$$H(t) \geq A_j(t+1)^{a_j}, G(t) \geq B_j(t+1)^{b_j}, j \in \mathbb{N}.$$

Under this assumption, the first inequality of (4.12) could be simplified as

$$\begin{aligned} H(t) &\geq e^{-\lambda(1-c)t} \int_{t/2}^t \frac{c_0 e^{\lambda(1-c)\tau} G^q(\tau)}{(\tau+1)^{(n-1)(p-1)/2}} d\tau + H(0) \\ &\geq \frac{c_1(G(t)/2)^p}{(t+1)^{(n-1)(p-1)/2}} \left(\frac{1 - e^{-\lambda(1-c)t}}{2\lambda(1-c)} \right) \geq \frac{c_2(G(t)^p}{(t+1)^{(n-1)(p-1)/2}}, \end{aligned}$$

for some constant $c_2(c_1, \lambda, p) > 0$ and $t \geq 2$. And the system (4.12) reads

$$(4.13) \quad \begin{cases} H \geq \frac{c_2|G|^p}{(t+1)^{(n-1)(p-1)/2}}, H(0) \geq c_2\varepsilon, \\ G'(t) \geq \frac{c_1H^q}{(t+1)^{(n-1)(q-1)/2}}, G(0) \geq c_1\varepsilon. \end{cases}$$

Then by the same way in case $c = 1$, there exists $\varepsilon_0 > 0$ such that for any $\varepsilon \in (0, \varepsilon_0)$, we have the iteration equation

$$(4.14) \quad \begin{cases} A_{j+1} = c_2 B_j^p, a_{j+1} = p b_j - \frac{(n-1)(p-1)}{2}, j \geq 1, \\ B_j = \frac{c_1 A_j^q}{b_j}, b_j = 1 + q a_j - \frac{(n-1)(q-1)}{2}, j \geq 1, \end{cases}$$

with

$$A_1 = c_2\varepsilon, a_1 = 0.$$

Hence we get that

$$\begin{aligned} a_j &= \left(\frac{p}{pq-1} - \frac{n-1}{2} \right) (pq)^{j-1} - \left(\frac{p}{pq-1} - \frac{n-1}{2} \right), j \in \mathbb{N}, \\ A_j &= c_2 B_{j-1}^p = c_1^p c_2 \frac{A_{j-1}^{pq}}{b_{j-1}^p} \geq K \frac{A_{j-1}^{pq}}{(pq)^{(j-1)p}}, K = c_1^p c_2 \left(\frac{p}{pq-1} - \frac{n-1}{2} + 1 \right)^{-p}. \end{aligned}$$

Then we see that

$$\begin{aligned} \ln A_j &\geq pq \ln A_{j-1} - (j-1)p \ln(pq) + \ln K \\ &\geq (pq)^2 \ln A_{j-2} - p \ln(pq) (pq(j-2) + (j-1)) + (pq+1) \ln K \\ &\geq (pq)^{j-1} \ln A_1 + (pq)^{j-1} \sum_{m=1}^{j-1} \frac{\ln K - mp \ln(pq)}{(pq)^m}. \end{aligned}$$

Note that the series

$$S(\infty) = \sum_{m=1}^{\infty} \frac{|\ln K| + mp \ln(pq)}{(pq)^m} < \infty,$$

thus we get that

$$\ln A_j \geq (pq)^{j-1} (\ln A_1 - S(\infty)).$$

Hence we have the lower bound estimate of $H(t)$

$$\begin{aligned} H(t) &\geq A_j(t+1)^{a_j} = e^{\ln A_j + a_j \ln(t+1)} \\ &\geq e^{(pq)^{j-1} \left(\ln A_1 - S(\infty) + \left(\frac{p}{pq-1} - \frac{n-1}{2} \right) \ln(t+1) \right)} e^{-\left(\frac{p}{pq-1} - \frac{n-1}{2} \right) \ln(t+1)}, \forall j \in \mathbb{N}, \end{aligned}$$

which yields the finite time blow up phenomenon if

$$\left(\ln A_1 - S(\infty) + \left(\frac{p}{pq-1} - \frac{n-1}{2} \right) \ln(t+1) \right) > 0,$$

so we get the lifespan estimate

$$T_\varepsilon \lesssim \varepsilon^{-\left(\frac{p}{pq-1} - \frac{n-1}{2} \right)^{-1}}.$$

5. PROOF OF THEOREM 1.3

Without loss of generality, we assume $0 < c \leq 1$. In this case, we see that $D_1 \subset D_2$.

5.1. Blow up for $\Gamma_{SG}(p, q, n) > 0$. In this case, we define functionals

$$F(t) = \int_{D_2} u_t dv_{\mathbf{g}}, \quad G(t) = \int_{D_2} v dv_{\mathbf{g}}.$$

Then by the system (1.19), we have

$$F'(t) = \int_{D_2} |v|^p dv_{\mathbf{g}}, \quad G''(t) = \int_{D_2} |u_t|^q dv_{\mathbf{g}}.$$

To connect F' , G'' with F , G , we apply Hölder's inequality to get the following ODE system

$$(5.1) \quad F' \geq c_0 \frac{G^p}{(t+1)^{n(p-1)}}, \quad G'' \geq c_1 \frac{F^q}{(t+1)^{n(q-1)}},$$

for some constants $c_0, c_1 > 0$. By the assumption (1.23) on the initial data, we have the lower bound of F , G

$$(5.2) \quad F(t) \gtrsim \varepsilon, \quad G \gtrsim \varepsilon(t+1), \quad t > 0.$$

By the same procedure in the proof of Theorem 1.1, the lower bound of F , G could be improved to

$$(5.3) \quad F(t) \gtrsim \varepsilon^p (t+1)^{n-\frac{n-1}{2}p}, \quad G(t) \gtrsim \varepsilon^q (t+1)^{n+1-\frac{n-1}{2}q}, \quad t > 0,$$

under the condition $n - \frac{n-1}{2}p > 0$, $n - \frac{n-1}{2}q > 0$, by auxiliary functions

$$H_1(t) = \int_{D_1} u_t \psi_1 dv_{\mathfrak{g}}, \quad H_2(t) = \int_{D_2} v \psi_2 dv_{\mathfrak{g}}.$$

In fact, by (1.19), we have

$$H_2'' + 2\lambda H_2' = \int |u_t|^q \psi_2 dv_{\mathfrak{g}} \geq 0,$$

thus by the assumption (1.23) on the initial data,

$$H_2'(t) \geq e^{-2\lambda t} H_2'(0) \geq 0, \quad H_2(t) \geq H_2(0) \gtrsim \varepsilon.$$

By applying Hölder's inequality and Lemma 2.2, we get

$$\begin{aligned} \varepsilon &\lesssim H_2 \lesssim \left(\int_{D_2} |v|^p dv_{\mathfrak{g}} \right)^{1/p} \left(\int_{D_2} |\psi_2|^{p'} dv_{\mathfrak{g}} \right)^{1/p'} \\ &\lesssim \left(\int_{D_2} |v|^p dv_{\mathfrak{g}} \right)^{1/p} \left((t+1)^{n-1-\frac{n-1}{2}p'} \right)^{1/p'}. \end{aligned}$$

Then

$$F' = \int_{D_2} |v|^p dv_{\mathfrak{g}} \gtrsim \varepsilon^p (t+1)^{n-1-\frac{n-1}{2}p},$$

thus we could integrate the above inequality get the lower bound of F in (5.3).

For the lower bound of auxiliary function $H_1(t)$, we need another functional $L(t) = \int_{D_1} (u_t \psi_1 + \lambda u \psi_1) dv_{\mathfrak{g}}$, which appeared in the proof of Theorem 1.2. It is easy to check that

$$(2H_1 - L)' + 2\lambda c(2H_1 - L) = \int_{D_1} |v|^p \psi_1 dv_{\mathfrak{g}} \geq 0, \quad L' = \int |v|^p \psi_1 dv_{\mathfrak{g}} \geq 0,$$

since $\text{supp } u \subset D_1$. Thus by the assumption on the initial data (1.23), we have

$$(2H_1 - L)(t) \geq (2H_1 - L)(0) \gtrsim \varepsilon, \quad L(0) \geq 0,$$

which yields

$$H_1(t) \geq \frac{L(t)}{2} + \frac{1}{2}(2H_1 - L)(0) \gtrsim \varepsilon.$$

By applying Hölder's inequality and Lemma 2.2, we get

$$\begin{aligned} \varepsilon &\lesssim H_1 \lesssim \left(\int_{D_1} |u_t|^q dv_{\mathfrak{g}} \right)^{1/q} \left(\int_{D_1} |\psi_1|^{q'} dv_{\mathfrak{g}} \right)^{1/q'} \\ &\lesssim \left(\int_{D_1} |u_t|^q dv_{\mathfrak{g}} \right)^{1/q} \left((t+1)^{n-1-\frac{n-1}{2}q'} \right)^{1/q'}, \end{aligned}$$

and

$$G'' = \int_{D_2} |u_t|^q dv_{\mathfrak{g}} \geq \int_{D_1} |u_t|^q dv_{\mathfrak{g}} \gtrsim \varepsilon^q (t+1)^{n-1-\frac{n-1}{2}q}.$$

Hence, we could integrate the above inequality twice to get (5.3).

Heuristically, the lower bound of F in (5.3) could be improved by the ODE system (5.1) if

$$\left(\left(n - \frac{n-1}{2}p \right) q - n(q-1) + 2 \right) p - n(p-1) + 1 > n - \frac{n-1}{2}p,$$

that is the region

$$(5.4) \quad \frac{2+p^{-1}}{pq-1} > \frac{n-1}{2} .$$

And the lower bound of G in (5.3) could be improved by (5.1) if

$$\left(\left(n+1 - \frac{n-1}{2}q \right) p - n(p-1) + 1 \right) q - n(q-1) + 2 > n+1 - \frac{n-1}{2}q ,$$

that is the region

$$(5.5) \quad \frac{p+1+q^{-1}}{pq-1} > \frac{n-1}{2} .$$

Combine (5.4) and (5.5), we could obtain the finite time blow up result on the range

$$\max\left\{ \frac{2+p^{-1}}{pq-1} , \frac{p+1+q^{-1}}{pq-1} \right\} - \frac{n-1}{2} > 0 .$$

In the following, we only give the the upper bound lifespan estimate when

$$\frac{p+1+q^{-1}}{pq-1} - \frac{n-1}{2} > 0 ,$$

since the other case follow the same way. In the same way as in the proof of Theorems 1.1-1.2, we have that there exists $\varepsilon_0 > 0$, such that for any $\varepsilon \in (0, \varepsilon_0)$, we could assume the lower bound of F, G by some sequences

$$F \geq A_j(t+1)^{a_j} , G \geq B_j(t+1)^{b_j} , j \in \mathbb{N} .$$

We start the iteration with

$$(5.6) \quad B_1 = \varepsilon^q, \quad b_1 = n+1 - \frac{n-1}{2}q ,$$

and by (5.1) we have the relation

$$(5.7) \quad \begin{cases} A_j = \frac{c_0 B_{j-1}^p}{a_j} , a_j = pb_{j-1} - n(p-1) + 1 , j \geq 2 \\ B_j = \frac{c_1 A_j^q}{(b_j-1)b_j} , b_j = qa_j - n(q-1) + 2 , j \geq 2 \end{cases}$$

Then we get that

$$b_j = \left(\frac{p+1+q^{-1}}{pq-1} - \frac{n-1}{2} \right) q(pq)^{j-1} + n - \frac{q+2}{pq-1} .$$

For the coefficient B_j , it is easy to see that

$$B_j = \frac{c_1 c_0^q B_{j-1}^{pq}}{(a_j)^q (b_j-1)b_j} \geq \frac{c_1 c_0^q B_{j-1}^{pq}}{(pb_j+1)^{q+2}} \geq \frac{M B_{j-1}^{pq}}{(pq)^{j(q+2)}}$$

where

$$M = c_1 c_0^q \left(\frac{p+1+q^{-1}}{pq-1} + \frac{q+2}{pq-1} + n+1 \right)^{-(q+2)} .$$

Hence we get

$$\begin{aligned}
\ln B_j &\geq (pq) \ln B_{j-1} + \ln M - j(q+2) \ln(pq) \\
&\geq (pq)^2 \ln B_{j-2} + (pq+1) \ln M - (pq(j-1) + j)(q+2) \ln(pq) \\
&\geq (pq)^{j-1} \ln B_1 - (pq)^{j-1} \sum_{k=1}^{j-1} \frac{(k+1)(q+2) \ln(pq) - \ln M}{(pq)^k}
\end{aligned}$$

Note that the series is convergence, that is

$$S(\infty) = \sum_{k=1}^{\infty} \frac{(k+1)(q+2) \ln(pq) + |\ln M|}{(pq)^k} < \infty .$$

Then we have

$$\ln B_j \geq (pq)^{j-1} (\ln B_1 - S(\infty)) .$$

Thus

$$\begin{aligned}
G &\geq B_j (t+1)^{b_j} \\
&\geq e^{(pq)^{j-1} \left(\ln B_1 - S(\infty) + \left(\frac{p+1+q^{-1}}{pq-1} - \frac{n-1}{2} \right) q \ln(t+1) \right)} e^{(n - \frac{q+2}{pq-1}) \ln(t+1)} , \forall j \in \mathbb{N} ,
\end{aligned}$$

which yields the lifespan estimate

$$T_{\varepsilon} \lesssim \varepsilon^{-\left(\frac{p+1+q^{-1}}{pq-1} - \frac{n-1}{2} \right)^{-1}} .$$

Similarly, if we assume

$$\frac{2+p^{-1}}{pq-1} - \frac{n-1}{2} > 0 ,$$

with the iteration relation

$$(5.8) \quad \begin{cases} A_j = \frac{c_0 B_j^p}{a_j} , a_j = pb_j - n(p-1) + 1 , j \geq 2 \\ B_j = \frac{c_1 A_{j-1}^q}{(b_j-1)b_j} , b_j = qa_{j-1} - n(q-1) + 2 , j \geq 2 \end{cases}$$

and

$$(5.9) \quad A_1 = \varepsilon^p , a_1 = n - \frac{n-1}{2} p ,$$

we will have that

$$T_{\varepsilon} \lesssim \varepsilon^{-\left(\frac{2+p^{-1}}{pq-1} - \frac{n-1}{2} \right)^{-1}} .$$

Hence, we obtain the lifespan estimate

$$T_{\varepsilon} \lesssim \min \left\{ \varepsilon^{-\left(\frac{p+1+q^{-1}}{pq-1} - \frac{n-1}{2} \right)^{-1}} , \varepsilon^{-\left(\frac{2+p^{-1}}{pq-1} - \frac{n-1}{2} \right)^{-1}} \right\} .$$

5.2. Blow up for $\Gamma_{GG}^*(p, q, n) > 0, c \neq 1$ and $\Gamma_{GG}(p, q, n) > 0, c = 1$. We define functionals

$$F(t) = \int_{D_2} (u_t \psi_1 + \lambda c u \psi_1) dv_{\mathfrak{g}} , H(t) = \int_{D_2} v \psi_2 dv_{\mathfrak{g}} .$$

Then by the system (1.10), we have

$$(5.10) \quad F'(t) = \int_{D_2} |v|^p \psi_1 dv_{\mathfrak{g}} , H''(t) + 2\lambda H' = \int_{D_2} |u_t|^q \psi_2 dv_{\mathfrak{g}} .$$

In order to relate F' with H , we let $H_1(t) = \int_{D_2} u_t \psi_1 dv_{\mathbf{g}}$ and it is easy to check that

$$(2H_1 - F)' + 2c\lambda(2H_1 - F) = \int_{D_2} |v|^p \psi_1 dv_{\mathbf{g}} \geq 0 ,$$

thus by the assumption on the initial data , we have

$$(2H_1 - F)(t) \geq (2H_1 - F)(0) \geq 0 ,$$

which yields

$$\frac{F}{2} \leq H_1(t) = \int_{D_2} u_t \psi_1 dv_{\mathbf{g}} .$$

By applying Hölder's inequality to H_1 and H , we have

$$\begin{aligned} \int_{D_2} |v|^p \psi_1 dv_{\mathbf{g}} &\gtrsim \frac{\left(\int_{D_2} v \psi_2 dv_{\mathbf{g}} \right)^p}{\left(\int_{D_2} \psi_1^{-\frac{p'}{p}} \psi_2^{p'} dv_{\mathbf{g}} \right)^{p/p'}} \gtrsim \frac{H^p}{\left(\int_{D_2} \psi_1^{-\frac{p'}{p}} \psi_2^{p'} dv_{\mathbf{g}} \right)^{p/p'}} , \\ \int_{D_1} |u_t|^q \psi_2 dv_{\mathbf{g}} &\gtrsim \frac{\left(\int_{D_1} u_t \psi_1 dv_{\mathbf{g}} \right)^q}{\left(\int_{D_1} \psi_2^{-\frac{q'}{q}} \psi_1^{q'} dv_{\mathbf{g}} \right)^{q/q'}} \gtrsim \frac{H_1^q}{\left(\int_{D_1} \psi_2^{-\frac{q'}{q}} \psi_1^{q'} dv_{\mathbf{g}} \right)^{q/q'}} . \end{aligned}$$

By (5.10) and Lemma 2.2, we get the following ODE system

$$(5.11) \quad \begin{cases} F'(t) \geq \frac{c_0 |H|^p}{e^{\lambda(c-1)t} (t+1)^{\frac{(n-1)(p-1)}{2}}} , & F(0) \geq c_0 \varepsilon , \\ H'' + 2\lambda H' \geq \frac{c_1 |F|^q}{e^{\lambda(1-c)t} (t+1)^{\frac{(n-1)(q-1)}{2}}} , & H(0) \geq c_1 \varepsilon , \end{cases}$$

for some constants $c_0, c_1 > 0$.

Note that the above system could be reduced to system (4.10). In fact, we have

$$(e^{2\lambda t} H')' = e^{2\lambda t} (H''(t) + 2\lambda H') \geq \frac{c_1 e^{2\lambda t} F^q}{e^{\lambda(1-c)t} (t+1)^{(n-1)(q-1)/2}} ,$$

thus

$$\begin{aligned} H' &\geq e^{-2\lambda t} \int_{t/2}^t \frac{c_1 e^{2\lambda \tau} F^q(\tau)}{e^{\lambda(1-c)\tau} (\tau+1)^{(n-1)(q-1)/2}} d\tau + H'(0) \\ &\geq \frac{c_1 (F(t/2))^q}{e^{\lambda(1-c)t} (t+1)^{(n-1)(q-1)/2}} \left(\frac{1 - e^{-\lambda t}}{2\lambda} \right) \\ &\geq \frac{c_2 (F(t/2))^q}{e^{\lambda(1-c)t} (t+1)^{(n-1)(q-1)/2}} , \quad t \geq 2 , \end{aligned}$$

for some constant $c_2 > 0$ depends on c_1 and λ . Hence we get the blow up result when $\Gamma_{GG}^*(p, q, n) > 0, c \neq 1$ and $\Gamma_{GG}(p, q, n) > 0, c = 1$ as well as the corresponding lifespan estimates.

REFERENCES

- [1] R. Agemi, Y. Kurokawa and H. Takamura, *Critical curve for p - q systems of nonlinear wave equations in three space dimensions*, J. Differential Equations, 167 (1) (2000) 87–133.
- [2] R. Agemi, K. Yokoyama, *The null conditions and global existence of solutions to systems of wave equations with different propagation speeds*, in “Advances in nonlinear partial differential equations and stochastics”, Series on Adv. in Math. for Appl. Sci., 48 (1998) 43–86.
- [3] W. Dai, D.Y. Fang, C.B. Wang, *Global existence and lifespan for semilinear wave equations with mixed nonlinear terms*, J. Differential Equations 267 (5) (2019) 3328–3354.
- [4] K. Deng, *Blow-up of solutions of some nonlinear hyperbolic systems*, Rocky Mountain J. Math., 29 (3) (1999) 807–820.
- [5] K. Hidano, K. Yokoyama, *Life span of small solutions to a system of wave equations*, Nonlinear Anal. 139 (2016) 106–130.
- [6] H. Kubo, K. Kubota, H. Sunagawa, *Large time behavior of solutions to semilinear systems of wave equations*, Math. Ann., 335 (2) (2006) 435–478.
- [7] A. Hoshiga, H. Kubo, *Global solvability for systems of nonlinear wave equations with multiple speeds in two space dimensions*, Differential Integral Equations, 17 (5-6) (2004) 593–622.
- [8] M. Ikeda, M. Sobajima, K. Wakasa, *Blow-up phenomena of semilinear wave equations and their weakly coupled systems*, Journal of Differential Equations, 267 (9) (2019) 5165–5201.
- [9] M. Kovalyov, *Resonance-type behavior in a system of nonlinear wave equations*, J. Differential Equations, 77 (1) (1989) 73–83.
- [10] H. Kubo, M. Ohta, *Critical blowup for systems of semilinear wave equations in low space dimensions*, J. Math. Anal. Appl., 240 (2) (1999) 340–360.
- [11] H. Kubo, M. Ohta, *Small data blowup for systems of semi-linear wave equations with different propagation speeds in three space dimensions*, J. Differential Equations. 163 (2) (2000) 475–492.
- [12] H. Kubo, M. Ohta, *On the global behavior of classical solutions to coupled systems of semilinear wave equations*, New trends in the theory of hyperbolic equations, 113–211, Oper. Theory Adv. Appl., 159, Adv. Partial Differ. Equ. (Basel), Birkhäuser, Basel, 2005.
- [13] N.A. Lai, H. Takamura, *Blow-up for semilinear damped wave equations with subcritical exponent in the scattering case*, Nonlinear Anal., 168 (2018) 222–237.
- [14] M.Y. Liu, C.B. Wang, *The blow up of solutions to semilinear wave equations on asymptotically Euclidean manifolds*, arXiv:1912.02540, 2019.
- [15] M.Y. Liu, C.B. Wang, *Blow up for small-amplitude semilinear wave equations with mixed nonlinearities on asymptotically Euclidean manifolds*, J. Differential Equations, 269 (10) (2020) 8573–8596.
- [16] D. Santo, *Global existence and blow-up for a hyperbolic system in three space dimensions*, Rend. Istit. Mat. Univ. Trieste, 29 (1-2) (1997) 115–140.
- [17] D. Santo, E. Mitidieri, *Blow-up of solutions of a hyperbolic system: the critical case*, (Russian) Differ. Uravn., 34 (9) (1998) 1155–1161, translation in Differential Equations, 34 (9) (1999) 1157–1163.
- [18] D. Santo, V. Georgiev, E. Mitidieri, *Global existence of the solutions and formation of singularities for a class of hyperbolic systems*, Geometrical optics and related topics, (1997) 117–140.
- [19] W. Xu, *Blowup for systems of semilinear wave equations with small initial data*, J. Partial Dif. Eqs, 17 (2004) 198–206.
- [20] K. Yokoyama, *Global existence of classical solutions to systems of wave equations with critical nonlinearity in three space dimensions*, J. Math. Soc. Japan, 52 (3) (2000) 609–632.

DEPARTMENT OF MATHEMATICS, ZHEJIANG SCI-TECH UNIVERSITY, HANGZHOU 310018, P. R. CHINA

Email address: mengyunliu@zstu.edu.cn