

J. Am. Chem. Soc., 1996, 118(26), 6141-6146, DOI:10.1021/ja960720z

## **Terms & Conditions**

Electronic Supporting Information files are available without a subscription to ACS Web Editions. The American Chemical Society holds a copyright ownership interest in any copyrightable Supporting Information. Files available from the ACS website may be downloaded for personal use only. Users are not otherwise permitted to reproduce, republish, redistribute, or sell any Supporting Information from the ACS website, either in whole or in part, in either machine-readable form or any other form without permission from the American Chemical Society. For permission to reproduce, republish and redistribute this material, requesters must process their own requests via the RightsLink permission system. Information about how to use the RightsLink permission system can be found at <a href="http://pubs.acs.org/page/copyright/permissions.html">http://pubs.acs.org/page/copyright/permissions.html</a>



## Supplementary Material

**Figure S1.** Characterization of the synthesis of ImPyPy-γ-PyPyPy-β-Dp. (a) (top) Analytical HPLC analysis of crude synthesis products from the machine-assisted solid phase synthesis, (a) (bottom) Analytical HPLC analysis of purified polyamide, C18 chromatography, 0.1% (wt/v) TFA, gradient elution 1.25% CH<sub>3</sub>CN/min monitored at 254 nm. (c) MALDI-TOF mass spectral analysis of purified polyamide.

